

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 February 2003 (20.02.2003)


PCT

(10) International Publication Number  
**WO 03/014361 A1**

- (51) International Patent Classification<sup>7</sup>: **C12N 15/67**, 15/64, 15/62, 5/10
- (74) Common Representative: **ALTANA PHARMA AG**; Byk-Gulden-Strasse 2, 78467 Konstanz (DE).
- (21) International Application Number: **PCT/EP02/08520**
- (81) Designated States (*national*): AE, AL, AU, BA, BR, CA, CN, CO, CU, DZ, EC, GE, HR, HU, ID, IL, IN, IS, JP, KR, LT, LV, MA, MK, MX, NO, NZ, PH, PL, RO, SG, SI, TN, UA, US, VN, YU, ZA, ZW.
- (22) International Filing Date: 31 July 2002 (31.07.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 01118632.7 2 August 2001 (02.08.2001) EP
- (84) Designated States (*regional*): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR).
- (71) Applicant (*for all designated States except US*): **ALTANA PHARMA AG** [DE/DE]; Byk-Gulden-Strasse 2, 78467 Konstanz (DE).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **HAUSER, Hans-jörg** [DE/DE]; Georg-Westermann-Allee 29, 38104 Braunschweig (DE). **MÜLLER, Peter, P.** [CH/DE]; Gerhart-Hauptmann-Strasse 8, 38239 Salzgitter-Thiede (DE). **SCHÄFER, Klaus, P.** [DE/DE]; Glockenbrunnenstrasse 6, 78465 Konstanz (DE). **STEINHILBER, Wolfram** [DE/DE]; Herzog-Erchanger-Strasse 16, 78333 Stockach (DE).
- Declarations under Rule 4.17:**
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AL, AU, BA, BR, CA, CN, CO, CU, DZ, EC, GE, HR, HU, ID, IL, IN, IS, JP, KR, LT, LV, MA, MK, MX, NO, NZ, PH, PL, RO, SG, SI, TN, UA, VN, YU, ZA, ZW, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR)
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only
- Published:**
- with international search report

[Continued on next page]

(54) Title: NOVEL RECOMBINANT GENE EXPRESSION METHOD BY STOP CODON SUPPRESSION



(57) Abstract: This invention describes a novel recombinant gene expression method based on a novel recombinant gene expression vector, comprising in the following order a promoter sequence, a gene of interest, a translational stop signal and translationally linked to said gene of interest a selectable marker gene.

WO 03/014361 A1



---

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## NOVEL RECOMBINANT GENE EXPRESSION METHOD BY STOP CODON SUPPRESSION

**Technical Field:**

The present invention relates to the field of recombinant protein expression in host cells, to a new method to obtain and identify cell clones expressing a gene of interest above a threshold level and stably maintaining or increasing expression levels during cultivation and to a process for producing recombinant proteins.

**Prior Art:**

The general principle of genetic modification of cultured cells or whole animals by various gene transfer methods is well known and a key strategy in both basic research and biotechnological applications. For large scale productions and also for many small scale applications efficient and stable expression of a recombinant product is an absolute requirement. However, difficulties exist in identifying host cells that produce recombinant protein at sufficient levels and in identifying host cell clones that maintain expression levels during prolonged cultivation periods.

Many approaches focusing on recombinant gene expression require co-expression of different gene products in a modified host cell. Frequently used procedures to generate a modified host cell producing a specific protein demand the introduction of foreign nucleic acid, e. g. an expression vector, into said host cell and the simultaneous expression of a selectable marker gene and a gene of interest. This is frequently achieved by co-transferring both genes on a separate DNA construct or alternatively by placing both gene expression cassettes on a single vector provided with two heterologous sets of transcriptional regulatory elements (e. g. pCDNA3 in Huang, C. F. et al., (2001) Disrupting the transforming activity of shrimp ras(Q61)K by deleting the CAAX box at the C-terminus. J. Exp. Zool. 289: 441 - 448). Frequently, the majority of clones that survive the selection process do not express a protein of interest above a threshold level. Usually below 1 % of selected clones are high level producers [Fussenegger, M. et al., (1999) Genetic optimization of recombinant glycoprotein production by mammalian cells. Trends in Biotechnology 17: 35 - 42]. Both approaches rarely result in the generation of cell lines that express the desired gene product over extended periods of time. Cell clones that express the product at moderate levels under selection pressure are frequently used for experimental purposes, but these cell clones do not meet industrial requirements. To obtain stable cell clones time consuming and laborious screening is necessary. The lack of expression stability using the above described gene expression methods is a consequence of the independent expression levels of a selectable marker gene and a gene of interest, i. e. a high selectable marker gene expression level in a given cell clone does not correlate with a high co-expression of said gene of interest. This situation

may be improved by using constructs in which both genes are transcribed from the same promoter, i.e. polycistronic expression constructs or bidirectional promoters.

EP-A-0117058 describes an expression vector capable of expressing a desired protein in a vertebrate host cell, which vector comprises a first DNA sequence encoding for a desired protein and a second DNA sequence encoding for a screening marker protein where both said DNA sequences are operationally linked to the same promoter sequence and separated by a translational stop and a start codon; said genes are not translationally linked. Consequently, both proteins are produced separately in mature form.

WO 8805466 describes compositions and methods for expressing gene products of interest in eukaryotic cells with a recombinant dicistronic DNA expression vector that comprises a gene of interest and a selectable marker gene wherein the sequences are physically linked to one another and depend on one common promoter. Thus a cell transfected by such a vector is capable of expressing the gene products from both of the genes.

WO 9708330 is related to new expression systems and in particular to an expression system in which a gene of interest is expressed at an optimal level. The invention provides a recombinant expression vector comprising a gene of interest and a selectable marker gene, wherein the selectable marker gene is arranged downstream of the gene of interest and a stop codon associated with the gene of interest is spaced from the start codon of said selectable marker gene at a distance which is sufficient to ensure that translation reinitiation is required before said selectable marker protein is expressed from corresponding mRNA. Both genes are not translationally linked.

Importantly, the above mentioned examples describe the presence of a translation initiation codon for the selection marker that is outside of the reading frame of the gene of interest. Increasing the number of in-frame stop codons terminating the gene of interest does not invariably lead to a reduced translation of the selection marker reading frame. The reading frames of the gene of interest and the selectable marker gene do not overlap and are not in a defined relationship.

WO0144516 describes a high throughput method of assaying for compounds that inhibit premature translation termination and nonsense mediated RNA decay in cells. The method described is based on the use of nucleic acid encoding a polypeptide, wherein the coding sequence for the polypeptide comprises a premature stop codon.

Zinoni et al. [Zinoni F et al. (1990) PNAS 87 : 4660] describe fdhF-lacZ fusion constructs to analyse the discrimination of the internal fdhF-mRNA UGA codon. According to table 2, fdhF-lacZ constructs comprise the following cassette: 5'-part of lacZ – fdhF + internal UGA – 3'-part of lacZ – stop codon.

Sogaard et al. [Sogaard TMM et al. (1999) Biochemistry 64: 1668] describe an in vivo system to monitor translation termination activity. The system is based on heat stable secreted alkaline

phosphatase (SEAP) where suppression at the stop codon tags said protein with an S-peptide. The fusion protein comprising S-peptide specifically binds to S-protein which is coated on microtiter plates. Comparison of SEAP activity in the supernatant with SEAP-activity coated on the microtiter plate is a mean to monitor in vivo translation termination.

Goldman et al. [Goldman E et al. (2000) FASEB 14: 603] describe the cloning of an unusual nucleotide sequence, H10, into a reporter system (using  $\beta$ -galactosidase) and the testing for expression in E.coli. Figure 1 discloses the experimental design. DNA constructs comprise in the following order a start for transcription, FLAG, H10 sequence comprising internal stop codons, an E-tag and a gene encoding  $\beta$ -galactosidase. Goldman et al. do not disclose a recombinant gene expression vector as disclosed herein and do also not describe a method to isolate stable expressing bacterial and mammalian cell clones.

Danielson et al. [Danielson S et al. (2001) Gene 272 : 267] describe the insertion of the wild type lipolase gene into a phagemid vector. The resulting construct encodes for the secreted expression of lipase, as fused to a 5 kDa serum albumin binding domain, followed by a suppressable TAG stop codon and a truncated version of the M13 phage coat protein 3. It also describes the expression of the lipase as a lipase-ABD fusion protein. Danielson et al. do not describe DNA constructs as disclosed herein and do not disclose a method to isolate stable expressing bacterial and mammalian cell clones.

Kollmus et al. [Kollmus H. *et al.*, (1996) Analysis of eukaryotic mRNA structures directing cotranslational incorporation of selenocysteine. Nucleic Acids Research 24:1195 - 1201] describe an assay system designated to facilitate analysis of DNA elements that influence stop codon suppression in mammalian cells. It is based on reporter genes encoding  $\beta$ -galactosidase and luciferase, which are fused in frame via a TGA stop codon. The DNA is transcribed under control of the SV40 promoter and translation leads to the synthesis of said reporter enzymes generating either  $\beta$ -galactosidase or, upon suppression of the TGA codon, a fusion protein of  $\beta$ -galactosidase and luciferase. Kollmus *et al.* do not describe a recombinant gene expression vector comprising a selectable marker gene, as specified according to this invention. In addition, Kollmus et al. do not disclose a method to isolate stable expressing bacterial and mammalian cell clones.

None of the above cited patent or literature documents does describe a process of producing a host cell line which cell line efficiently expresses a gene of interest.

In summary, even with polycistronic constructs, high selection pressure or prolonged cultivation of cells result in the isolation of cell clones that efficiently express the selectable marker gene but not the gene of interest. Therefore, it is a considerable problem that there is no reliable and for the average research laboratory affordable method to isolate stable expressing bacterial and mammalian cell clones that express a gene of interest at a high level.

**Description of the Invention:**

This invention describes a novel recombinant gene expression method based on a novel recombinant gene expression vector comprising a gene encoding a selectable marker protein which is separated by a translational stop signal from an upstream arranged gene of interest, whereby both genes are translationally linked. Consequently, the expression of said selectable marker gene may be reduced compared to the expression rate of said gene of interest.

Surprisingly, it has now been found that a high proportion of expressing single cell clones may be obtained under selection pressure by the usage of said novel recombinant gene expression vector according to this invention. In said expressing single cell clones the stop codon-dependent translational coupling of marker gene expression with gene of interest expression results in two recombinant gene products, i. e. a product encoded by the gene of interest and as a fusion protein the gene of interest combined with the selectable marker gene.

More surprisingly, the novel recombinant gene expression vector can be used to achieve stable gene expression under selection conditions by resisting cellular mechanisms that uncouple the resistance to selection conditions from the expression of a recombinant gene of interest. Thus, the expression of the gene of interest is enhanced or stabilized by selection for activity of the encoded selectable marker protein.

In particular surprisingly, it was also found that stop codon suppression could be applied to reach selection marker gene expression levels and selection marker activity sufficient to achieve resistance of transfected cells to the selection procedure. It is even more particularly surprising that said cell clones comprising said novel recombinant gene expression vector and expressing said selection marker gene at levels sufficient to achieve resistance required for cell survival under selective conditions express said gene of interest above a threshold level, whereby said level is at least identical or superior to the yields generated by cell clones comprising a standard recombinant gene expression vector and expressing a gene of interest based on standard techniques.

It is surprising that a host cell comprising a recombinant gene expression vector as specified herein comprising both a translational stop signal and a gene of interest encoding a secreted product protein is capable in producing a homogenous product of said gene of interest whereby the amount of homogenous gene product produced is identical or superior to the yields generated by host cells comprising a standard recombinant gene expression vector comprising a gene encoding a secreted product protein.

The advantage of this system is an extremely tight coupling of selectable marker gene expression and gene of interest expression. This property results in several advantageous features as follows:

- Increased frequency of transformed/transfected host cells expressing said gene of interest after selection for cells expressing the selectable marker gene.
- By limiting the expression of the selectable marker gene at the translational level, transformed/transfected host cells with an increased transcriptional activity of the recombinant genes are selected. Thus the average recombinant protein expression of the gene of interest is also increased when compared to expression vectors without limiting selection gene expression.
- Increased stability of gene of interest expression under selective conditions when compared to conventional expression vectors with other mechanisms of selection, such as polycistronic vectors [Mueller, P. P. et al. (1999) Recombinant glycoprotein product quality in proliferation controlled BHK-21 cells. *Biotech. and Bioeng.* 65: 529 - 536] or vectors with bidirectional promoters [Baron, U. et al. (1995) Co-regulation of two gene activities by tetracycline via a bidirectional promoter. *Nucleic Acids Research* 23: 3605 - 3606].

In a first embodiment of the invention there is provided a recombinant gene expression vector comprising in the following order a promoter sequence, a gene of interest, a translational stop signal and translationally linked to said gene of interest a selectable marker gene, whereby a recombinant gene expression vector is excluded which recombinant gene expression vector contains in the following order a promoter sequence, a gene encoding  $\beta$ -galactosidase, a TGA stop codon, and translationally linked to said  $\beta$ -galactosidase gene a gene encoding luciferase.

In a second embodiment of this invention there is provided a recombinant gene expression vector as disclosed before comprising more than one gene of interest.

Preferably, the present invention provides a recombinant gene expression vector comprising said elements in said order whereby the gene of interest and the selectable marker gene are translationally linked. In accordance to this invention, translationally linked refers to a reading frame fusion, such that ribosomes translate the selection marker from the same translation initiation site as the gene of interest. It is particularly preferred that translationally linked refers to constructs wherein the gene of interest and the selectable marker gene are entirely or partially in the same reading frame. The gene of interest and the selectable marker gene may be separated either by a translational stop signal and/or translational frameshifting signals.

This invention refers to any sequence of desoxyribonucleotides which result in (1) an in frame arrangement of said gene of interest and said selectable marker gene and (2) a mRNA product comprising both said gene of interest and said selectable marker gene. Thereby, the gene of interest is located at the 5'-end of the reading frame fusion and it may be translated independently of the downstream arranged selectable marker gene. In addition, it is an arrangement whereby the reading frame of said selectable marker gene is arranged downstream of said gene of interest in a way that the translation of the selectable marker gene reading frame is dependent on prior translation of the gene of

interest such that the selectable marker gene product may be expressed as a fusion protein. In particular, the invention relates to said recombinant gene expression vector in a way that a separate translational start codon for the selectable marker gene reading frame may be omitted.

According to this invention, a gene refers to any sequence of deoxyribonucleotides encoding the information for the synthesis of a primary RNA transcript or a protein. A gene of interest refers to any nucleic acid sequence used for cloning in said recombinant gene expression vector encoding a protein or protein fragment thereof whose expression is of interest. According to this invention, a gene of interest refers to a wildtype or a modified gene, whereby a wild-type gene refers to any gene that predominates in the population because it confers the greatest fitness (ability to survive and reproduce) and whereby a modified gene refers to any gene which sequence differs from that of the wildtype gene either by mutation, deletion, insertion of a single or multiple nucleotide sequences or by fusion of additional sequence to at least one end of the gene sequence. In a preferred embodiment of this invention, a gene of interest refers to a gene encoding a protein of pharmaceutical interest such as (1) biotechnological products such as cytokines (Insulin, EPO (erythropoietin), TPO (Tissue Plasminogen Activator), GCSF (Granulocyte Colony Stimulating Factor), GM-CSF (Granulocyte-Macrophage Colony Stimulating Factor), blood clotting factors FVIII or FVII), (2) other pharmaceutically active products such as surfactant protein (SP) selected from the group of SP-A, SP-B, SP-C or SP-D, (3) gene products used for identification of drug targets, (4) any EST (Expressed Sequence Tag) reading frame, and (5) any reporter gene, in particular SEAP (Secreted Alkaline Phosphatase), *lacZ* ( $\beta$ -Galactosidase), *luc* (firefly or renilla luciferase), GFP (Green Fluorescence Protein) and derivatives thereof and proteins with similar function such as Red Fluorescence Protein (RFP), CAT (Chloramphenicol Acetyl Transferase), proteins and reporter protein tags exposed at the cell surface such as myc tag, polyhistidine tag, or flag antigen. In a particularly preferred embodiment of the present invention, said gene of interest preferably encodes a pharmaceutically active protein. A pharmaceutically active protein is understood as any protein capable for usage in diagnosis, prophylaxis, treatment or modification of physiological functions in human or animal body.

In the present invention, one promoter controls the expression of at least one gene of interest and a selectable marker gene. A promoter refers to any DNA sequences involved in promoting transcription of a nucleic acid sequence. In particular, a promoter refers to LTR (Long Terminal Repeat) promoter elements of Retrovirus such as MMTV (Mouse Mammary Tumor Virus), RSV (Rous Sarcoma Virus), MPSV (Myelo-Proliferative Sarcoma Virus). Particularly, a promoter refers to other viral promoters such as SV40 (Simian Virus) early or late promoter, or CMV (CytoMegalovirus) promoter. It also refers to the EF (Elongation Factor) promoter or to an artificial promoter such as tetracycline regulated promoter [Gossen, M. and Bujard, H. (1992) Tight control of gene expression in mammalian cells by tetracycline-responsive promoters. PNAS 89: 5547 - 5551], or to any other known promoter, such as eukaryotic, prokaryotic or viral promoters.

The vector used in the process of the invention may be any of the known types, comprising any DNA or RNA segment that can replicate autonomously within a cell, in particular in a bacterial cell, e. g. ex-



pression plasmid or viral vectors. In another embodiment of this invention, a vector refers to any DNA segment that can not replicate autonomously within a cell and which DNA segment integrates into the chromosome of said cell.

In a third embodiment of the invention, there is provided a recombinant gene expression vector comprising in the following order a promoter sequence, a gene of interest, a translational stop signal and translationally linked to said gene of interest a selectable marker gene encoding a functional protein, whereby a recombinant gene expression vector is excluded which recombinant gene expression vector contains in the following order a promoter sequence, a gene encoding  $\beta$ -galactosidase, a TGA stop codon, and translationally linked to said  $\beta$ -galactosidase gene a gene encoding luciferase.

Usually, bacterial cells transformed with a plasmid vector are positively selected for dominant antibiotic resistance markers carried by said plasmid, effectively maintaining a population of plasmid-containing cells. The same or alternative markers are used in transfected eukaryotic systems, e. g. neomycin phosphotransferase. Other selection or screening markers can be used if the activity of the fusion gene is high enough to allow marker activity above background levels.

This invention also refers to a recombinant gene expression vector, wherein said selectable marker gene for selection of transfected cells is a drug resistance gene encoding a protein that confers resistance to selection conditions, e. g. neo (neomycin phosphotransferase), hyg (hygromycin acetyltransferase), tk (Herpes simplex thymidine kinase), PAC (pyromycin acetyltransferase), zeo (zeozin resistance gene), DHFR (dihydrofolate reductase).

In a variant of this invention, said selectable marker gene refers to a reporter gene, in particular to GFP (Green Fluorescence Protein), whose expression can be used to differentiate between host cells producing said selectable marker gene and those which do not express said gene.

In particular, the invention refers to a recombinant gene expression vector comprising a selectable marker gene encoding neomycin phosphotransferase. A transfected host cell line comprising this vector may be positively selected from a plethora of cells lacking said vector by using G418. G418 is an aminoglycoside antibiotic that bind to the ribosome and reduce the fidelity of translation. G418 distinguishes between prokaryotic and eukaryotic ribosomes. G418 is a substrate for neomycin phosphotransferase encoded by the bacterial *neo* gene, resulting in the phosphorylation and thereby the inactivation of G418 [Beck, E. et al. (1982) Nucleotide sequence and exact localization of the neomycin phosphotransferase gene from transposon Tn5. *Gene* 19: 327 - 336; Colbere-Garapin, F. et al. (1981) A new dominant hybrid selective marker for higher eukaryotic cells *J. Mol. Biol.* 150: 1 - 13].

In a fourth embodiment of this invention there is provided a recombinant gene expression vector as disclosed before, wherein said translational stop signal is at least one stop codon selected from the group of TAA, TGA and TAG.

According to this invention, said selectable marker gene is arranged downstream of said gene of interest, whereby both genes are separated by a translational stop signal. A translational stop signal refers to any genetically element suitable to terminate the translation of a reading frame and thereby decrease the rate of translation of any reading frame encoded in the mRNA downstream of said translational stop signal. In particular, a translational stop signal refers to at least one stop codon on mRNA selected from the group of UGA, UAA and UAG, encoded by TGA, TAA and TAG, respectively, on the corresponding DNA construct. In another embodiment of this invention, more than one stop codon, preferably two or three similar or different stop codons are selected from said group. Consequently, the expression of said selectable marker gene is reduced when compared to a conventional translation fusion whereby the stop codon of the upstream gene is removed. In the absence of additional genetic elements, the reduction of the selection marker gene expression is defined as the natural rate of stop codon read-through. In mammalian cells, the naturally occurring stop codon read-through efficiency is about 0,1% [Kollmus, H. *et al.* (1996) Analysis of eukaryotic mRNA structures directing cotranslational incorporation of selenocysteine. *Nucleic Acids Research* 24:1195 - 1201]. By using several stop codons in frame the naturally stop codon-read-through efficiency can be further reduced or enhanced, respectively.

The read-through efficiency may be modulated by including a stop codon suppression mechanism. This can be a SECIS element for UGA suppression, suppressor tRNAs, ectopic expression of wild-type tRNAs, use of alternative stop codons UAG or UAA, by changing the context of the stop codon, by introducing a frameshift, or by the addition of an alternative transcriptional stop signal. An alternative transcriptional stop signal can be a natural cleavage and poly(A) addition site or a poly(A) consensus sequence inserted preferably downstream of the stop codon of the gene of interest, or in the 5'-region of the selection marker gene [Minvielle-Sebastia, L., Keller, W. (1999) mRNA polyadenylation and its coupling to other RNA processing reactions and to transcription. *Curr. Opin. Cell. Biol.* 11: 352 - 357; Wahle, E., Ruegsegger, U. (1999) 3'-End processing of pre-mRNA in eukaryotes. *FEMS Microbiol Rev* 23: 277 - 295; Edwalds-Gilbert, G., Veraldi, K. L., Milcarek, C. (1997) Alternative poly(A) site selection in complex transcription units: means to an end? *Nucleic Acids Research* 25: 2547 - 2561].

In a fifth embodiment of this invention, there is provided a recombinant gene expression vector comprising a promoter sequence, a gene of interest, a translational stop signal, a selectable marker gene (arranged in frame) and a SECIS element. The SECIS element (selenocystein insertion sequence) is a structural element (hairpin structure) in eukaryotic mRNA that enables the insertion of the 21<sup>st</sup> amino acid, selenocystein, at sites of the stop codon UGA [Kollmus H *et al.* (1996) Analysis of eukaryotic mRNA structures directing cotranslational incorporation of selenocysteine. *Nucleic Acids Research* 24: 1195 - 1201; Fagegaltier, D. *et al.* (2000) Structural analysis of new local features in SECIS RNA hairpins. *Nucleic Acids Research* 28: 2679 - 2689] and therefore increases the stop codon-read-through and the expression of any downstream localized genes.

Kollmus *et al.* measured the reduction in stop codon suppression efficiency in BHK-21 cells using the pig heart phospholipid hydroperoxidase glutathione peroxidase (PHGPx) gene. This SECIS element

supported a stop codon suppression activity of 2,8 %. A minimum SECIS element supported a 10-fold reduced activity of 0,3 % stop codon-read-through [Kollmus, H. et al. (1996) Analysis of eukaryotic mRNA structures directing cotranslational incorporation of selenocysteine. *Nucleic Acids Research* 24: 1195 - 1201]. The SECIS element of the gene encoding the rat 5'-deiodinase results in a relative stop codon suppression efficacy of 1,1% [Kollmus H et al. (1996) Analysis of eukaryotic mRNA structures directing cotranslational incorporation of selenocysteine. *Nucleic Acids Research* 24: 1195 - 1201]. A mutation in this element reduces the efficiency to 0,16%.

According to Kollmus et al. [Kollmus, H. et al. (1996) Analysis of eukaryotic mRNA structures directing cotranslational incorporation of selenocysteine. *Nucleic Acids Research* 24: 1195 - 1201], the stop codon suppression efficiency can be calculated by relating expression activity of luciferase - determined by measuring the enzymatic activities of luciferase - of cells containing certain bacterial plasmids to values obtained with a control plasmid. Based on this enzymatic method to determine the stop codon suppression efficiency, a stop codon suppression efficiency of 0,01 % to 10 % is a preferred embodiment of this invention. In particular, this invention refers to a stop codon suppression efficiency of 0,1 % to 5 %.

In said recombinant expression vectors, said gene of interest and said selectable marker gene are translationally linked. Therefore, a read-through results in the production of a fusion protein comprising both products of said gene of interest and said selectable marker gene. The function of a protein encoded by said selectable marker gene is conserved within the fusion protein, as exemplified by its ability demonstrated herein to confer antibiotic resistance to cells transfected with these constructs (see Figure 3). In that way, host cells comprising said recombinant expression vector may be selected by positive selection on expression of said fusion protein.

In a further embodiment of this invention, there is provided a host cell comprising a recombinant gene expression vector as disclosed herein which host cell is capable of expressing both said gene of interest and as a fusion protein said gene of interest and said selectable marker gene.

According to this invention, a host cell comprising a recombinant gene expression vector as disclosed herein comprises a eukaryotic or a prokaryotic cell line transfected/transformed with said recombinant gene expression vector.

According to this invention, transformation is synonymous with transfection.

According to this invention, a eukaryotic host cell line comprising said recombinant gene expression vector refers to any type of cell comprising a cell culture cell such as BHK-21, CHO (various Chinese Hamster Ovary cell lines), 293 (human kidney carcinoma cell line), NIH 3T3 (murine cell lines), SP2/0 (hybridoma), other eukaryotic cell lines of pharmaceutical or laboratory research interest, primary cells such as ES cells (Embryonic Stem cells), other mammalian stem or progenitor cells such as bone marrow progenitor cells, putting together a whole organism, whereby said recombinant gene

expression vector has been transferred into said host cell line by transfection. According to this invention, transfection refers to (1) the techniques specified in Sambrook *et al.* [Sambrook *et al.* (1989) In: Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, second edition], e.g. calciumphosphate-transfection, transfection using DEAE (Diethylaminoethyl)-dextran, lipofection, (2) electroporation and (3) infection, e.g. via a viral vector system. In another variant of this invention, said process of transfecting a host cell line comprises specific techniques. In particular, specific techniques refer to techniques revealed by following publications: Wigler, M. *et al.* (1977) Transfer of purified herpes virus thymidine kinase gene to cultured mouse cells. *Cell* 11: 223 - 232; Chesnoy S and Huang L (2000) Structure and function of lipid-DNA complexes for gene delivery. *Annu. Rev. Biophys. Biomol. Struct.* 29: 27 - 47; De Smedt SC, Demeester, J., Hennink, W. E., (2000) Cationic polymer based gene delivery systems. *Pharm. Research* 17: 113 -26; Mahato, R. I. (1999) Non-viral peptide-based approaches to gene delivery. *J. Drug Target* 7: 249 - 268; Stone, D. *et al.* (2000) Viral vectors for gene delivery and gene therapy within the endocrine system. *J. Endocrinol* 164: 103 - 118; Zhao, X., (2000) Gene transfer and drug delivery by electronic pulse delivery. A nonviral delivery approach. *Methods Mol. Biol.* 133: 37 - 43. As an alternative, other methods not known to a person skilled in the art can be used. For selection of transfected cells, methods described in Sambrook *et al.* [Sambrook *et al.* (1989) In: Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, second edition], or methods specified in the above mentioned publications concerning transfection of eukaryotic cells and other selection methods known to a person skilled in the art can be used.

In the case of a bacterial host cell the transformation and selection of positive clones may be performed according to the techniques specified in Sambrook *et al.* [Sambrook *et al.* (1989) In: Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, second edition] or by electroporation, bacteriophage infection or other methods known to a person skilled in the art.

Said transformation methods may be used by using said expression vector and selecting a transformed/transfected host cell on the product expression of said selectable marker gene. This process refers to both prokaryotic and eukaryotic cells. More preferred, this process refers to cells that are unstable or unpredictable in the expression level of recombinant proteins.

In a further embodiment of this invention, there is provided a process for producing a host cell clone comprising a recombinant gene expression vector as disclosed herein, which process comprises transforming a host cell with said recombinant gene expression vector and selecting a host cell clone comprising said recombinant gene expression vector on the expression of said selectable marker gene.

According to this invention, a host cell clone refers to a group of host cells, all of which are derived from a single individual cell through asexual reproduction. Except from changes that come about by mutation, all members of a host cell clone are genetically identical. Continued growth of a host cell clone in a laboratory culture results in a host cell line.

According to this invention a transformed/transfected host cell comprising said recombinant gene expression vector particularly express a sufficient amount of protein encoded by said selectable marker gene being an adequate amount of said selectable marker gene product to result in cell clones that survive the selection conditions after transformation of host cells with said recombinant expression vector under conditions that lead to cell death of comparable cells which do not carry said recombinant gene expression vector. In particular, a sufficient amount of protein encoded by said selectable marker gene refers to an adequate amount of said selectable marker gene product to select at least one transformed host cell out of a plethora of cells after introducing DNA or RNA of said recombinant expression vector into host cells.

In another embodiment of this invention there is provided the use of a recombinant gene expression vector as disclosed herein in a process for efficiently selecting a host cell clone highly expressing said gene of interest.

According to this invention, efficiently selecting refers to a method of transforming host cells, selecting said transformed host cells on the expression of said selectable marker gene – whereby those host cells survive the selection conditions which carry said recombinant expression vector and express said selectable marker gene under conditions that lead to cell death of comparable cells which do not carry said recombinant gene expression vector - and identifying at least 5% of host cell clones highly expressing said gene of interest compared to the total number of host cell clones surviving selection conditions. In particular, efficiently selecting refers to a selection method as mentioned before, whereby at least 10% of host cell clones highly expressing said gene of interest survive selection conditions. It is particularly preferred that efficiently selecting refers to a selection method as mentioned before, whereby at least 14% of host cell clones highly expressing said gene of interest survive selection conditions.

According to this invention, highly expressing said gene of interest refers to expressing a protein encoded by said gene of interest above a threshold level. In the meaning of this invention, a threshold level refers to at least 0,5 % of total cell protein. In particular, a threshold refers to at least 5 % of total cell protein. It is preferred, that a threshold level refers to at least 25 % of total cell protein. It is particularly preferred that a threshold level refers to at least 50 % of total cell protein. It is even more preferred that a threshold level refers to at least 75 % of total cell protein. In particular, a threshold level of said recombinant expressed protein produced in mammalian cells refers to at least 1  $\mu$ g protein per 1 ml culture supernatant final concentration. For mammalian cells a threshold level of at least 10  $\mu$ g/ml is preferred. It is particularly preferred that said threshold level refers to at least 25  $\mu$ g/ml. It is even more particularly preferred that said threshold level refers to at least 50  $\mu$ g/ml. Specifically, it is even more particularly preferred that said threshold level refers to at least 100  $\mu$ g/ml.

In accordance to this invention, protein concentrations are determined according to the methods described by Sambrook et al. [Sambrook et al. (1989) In: Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, second edition].

In a further embodiment of this invention there is also provided a process of producing a host cell line highly expressing a gene of interest, which process comprises a process for producing a host cell clone as disclosed herein comprising transforming a host cell with said recombinant gene expression vector, selecting a host cell clone comprising said gene expression vector on the expression of said selectable marker gene and identifying a host cell clone which highly expresses said gene of interest.

A transformed/transfected host cell comprising said recombinant expression vector is capable of expressing a sufficient amount of protein encoded by said selectable marker gene being an adequate amount of said selectable marker gene product to result in cell clones that survive the selection conditions after transformation of host cells with said recombinant expression vector under conditions that lead to cell death of comparable cells which do not carry said recombinant gene expression vector. In particular, a sufficient amount of protein encoded by said selectable marker gene refers to an adequate amount of said selectable marker gene product to select at least one transformed host cell out of a plethora of cells after introducing DNA or RNA of said recombinant expression vector into host cells.

In a further embodiment of this invention, there is provided a process for producing a protein encoded by a gene of interest, which process comprises the production of a host cell line comprising a recombinant gene expression vector, the cultivation of said host cell line, stably maintaining the expression of said gene of interest, and recovery of the product of said gene of interest, whereby said recombinant gene expression vector comprises in the following order a promoter sequence, a gene of interest, a translational stop signal and translationally linked to said gene of interest a selectable marker gene, and whereby the production of said host cell line comprises the transformation with said recombinant gene expression vector and the selection of a host cell comprising said recombinant gene expression vector on the expression of said selectable marker gene.

According to this invention, stably maintaining the expression of said gene of interest during cultivation refers to approximately constant production of proteins encoded by said gene of interest and said selectable marker gene over the period of 10 passages of the cell line. In particular, 25 cell passages are preferred. It is particularly preferred, that said host cell line produces said proteins at approximately constant level over 50 cell cycles. It is even more particularly preferred, that said host cell line produces said proteins at approximately constant level over 100 cell cycles. In a variant, it is preferred that stably maintaining the expression of said gene of interest during cultivation of transformed bacterial cells refers to recombinant protein productivity in bacteria that are superior to the stability of productivity obtained with related DNA constructs that do not employ the gene fusion technology described herein.

In another embodiment of this invention there is provided a host cell line comprising a recombinant gene expression vector as disclosed herein which host cell line is stably maintaining the expressing of said gene of interest during cultivation.

Most eukaryotic proteins are synthesized in the cytoplasm. Therefore, proteins must carry recognizable sequences or structures which allow them to be transported to the appropriate cellular compartment. Preproteins contain conserved amino acid residues, particularly found at the termini of preproteins (signal peptides) containing the information for cellular transporting. In eukaryotes, proteins destined for secretion, i.e. secreted product proteins, are initially targeted to the endoplasmatic reticulum (ER). This requires a N-terminal hydrophobic sequence signal peptide. The signal sequence is cleaved as the polypeptide is cotranslationally entering the ER lumen. Transmembrane proteins, e.g. placental alkaline phosphatase and many cell surface receptor subunits such as the platelet derived growth factor (PDGF) receptor, possess an internal hydrophobic stop transfer sequence often flanked with positively charged residues [Dalbey, R. E. (1990) Positively charged residues are important determinants of membrane protein topology. Trends Biochem. Sci. 15: 253 - 257]. The presence of a membrane-spanning domain in a secreted protein retains such a protein in cellular membranes such as the ER membrane, Golgi or cell membrane. Removal of the membrane-spanning domain in such proteins results in the secretion, as it is the case with SEAP, the secreted form of alkaline phosphatase. Membrane proteins of bacterial cells may also have hydrophobic stop transfer sequences to ensure their incorporation into the bacterial membrane.

In a further embodiment, there is also provided a recombinant gene expression vector comprising in the following order a promoter sequence, a gene of interest, a translational stop signal, a stop transfer sequence and translationally linked to the gene of interest a selectable marker gene. Said expression system is applicable to secreted product proteins.

Eukaryotic host cells comprising said recombinant gene expression vector may be positively selected on the expression of said fusion protein, comprising said selectable marker. Fusion proteins may be retained in the ER lumen thereby mediating resistance, whereas a product of said gene of interest is secreted in the cell's surrounding culture medium. This system allows a simple and selective recombinant production of a secreted product protein.

In another embodiment of this invention, there is provided a process for producing a secreted product protein encoded by a gene of interest, which process comprises the production of a host cell line comprising a recombinant gene expression vector, the cultivation of said host cell line, stably maintaining the expression of said gene of interest, and recovery of said secreted product protein from the cell's surrounding culture medium, whereby said recombinant gene expression vector comprises in the following order a promoter sequence, a gene of interest encoding a secreted product protein, a translational stop signal, an in frame stop transfer sequence and translationally linked to said gene of interest a selectable marker gene, and whereby the production of said host cell line comprises the transformation with said recombinant gene expression vector and the selection of a host cell comprising said recombinant gene expression vector on the expression of said selectable marker gene. In a variant of this invention said recombinant gene expression vector comprising said genetic element may also be used for expression of secreted product proteins in bacterial cells.

The invention will now be particularly described by way of examples with reference to the given figures. The following examples do not limit the scope of the invention.

### **Description of Diagrams:**

**Fig. 1. Expression constructs used for stop codon suppression dependent expression.**

Plasmids were constructed from DNA fragments containing transcriptional units encoding the drug resistance gene neomycin phosphotransferase (*neo*) or the GFP-*neo* fusion protein (GFP*neo*). Plasmids were constructed with a GFP-*neo* fusion protein separated by an in-frame stop codon (GFP*stopneo*), with a GFP-*neo* fusion protein separated by an in-frame stop codon and followed by an element (SECIS) that promotes the insertion of the amino acid selenocystein at the stop codon (GFP*stopneo*-SECIS), and with a secreted alkaline phosphatase (SEAP) reading frame terminated by a stop codon. The downstream sequences can only be translated when an error occurs and the ribosomes do not terminate translation at the stop codon. SEAP is followed by a membrane spanning domain (M) fused to the *neo* reading frame (SEAP*stopMneo*); the membrane spanning domain is derived from the platelet-derived growth factor receptor (PDGFR) isolated from the plasmid pDISPLAY (Invitrogen) and serves the purpose to anchor the secreted protein to the plasma membrane. Thereby the secreted product SEAP is exposed on the extracellular side, while neo protein is on the intracellular face of the membrane. The intracellular localization is considered a prerequisite to confer antibiotic resistance.

**Fig. 2. Translation products encoded by the GFP*neo* and GFP*stopneo* reading frames.**

The GFP*neo* reading frame (A, square box) encodes a single GFP*neo* fusion protein (A, upper arrow). The GFP*stopneo* construct contains the GFP reading frame (that is terminated by an in-frame stop codon) and the *neo* reading frame. The GFP reading frame encodes the GFP protein (B, upper arrow) and the natural rate of stop codon suppression would result in a small amount of GFP-*neo* fusion protein (B, lower arrow).

**Fig. 3. Frequency of resistant cell clones obtained with stop codon suppression dependent expression constructs.**

Equal amounts of plasmid DNA encoding the drug resistance gene *neo* (*neo*; GFP-*neo* fusion (GFP*neo*); a GFP-*neo* fusion separated by an in-frame stop codon (GFP*stopneo*); a GFP-*neo* fusion separated by an in-frame stop codon followed by a stop codon suppressing element (GFP*stopneo*-SECIS) were used to stably transfect BHK-21 cells (*neo*) using the calcium phosphate coprecipitation method (Pellicer, A. et al. (1978) The transfer and stable integration of the HSV thymidine kinase gene into mouse cells. Cell 14: 133 - 141.). After selection of surviving G418 resistant cells in the presence



of 1000  $\mu\text{g/ml}$  of the drug G418 in standard DMEM (Gibco) with 10 % fetal calf serum albumin (FCS) at a temperature of culture of 37°C in a 5 % CO<sub>2</sub> incubator, the number of clones was determined for each construct and compared to the number of clones obtained with a separate *neo* construct (Frequency of resistant cell clones). The bar on top of the columns indicates the standard deviation.

**Fig. 4. Gene expression in cell clones with stop codon suppression dependent translation.**

BHK-21 cells were stably transfected with the plasmid DNA indicated (see Figure 1). After selection for *neo* expressing clones in G418 containing cell culture medium, GFP fluorescence intensity of single suspended cells was determined by Fluorescence Activated Cell Sorter (FACS) analysis.

**Fig. 5. Stability of expression during cultivation under stop codon suppression dependent selection conditions.**

Stability of GFP expression from a GFP-*neo* reading frame fusion construct separated by an in-frame stop codon (GFP<sup>stopneo</sup>) was examined during extended cultivation time in 3 different single cell clones relative to a non-fluorescent reference cell clone by FACS analysis of the fluorescence from individual cells. Shown are the percentage of cells with GFP fluorescence above non-GFP-containing cell fluorescence intensity.

**Fig. 6. Secreted protein expression from stop codon suppression dependent translation constructs in transient transfection assays without selection.**

BHK-21 cells were transiently transfected with an IgG expression construct [Geserick C *et al.* (2000). Enhanced productivity during controlled proliferation of BHK-21 cells in continuously perfused bioreactors. Biotech and Bioeng 69: 266-274] and the SEAP expression constructs indicated by using the calcium coprecipitation method. IgG ELISA and SEAP activities from cell culture supernatants were determined 2 days after transfection. The SEAP activity is given relative to the expression level of the original SEAP construct and an IgG standard [Geserick C *et al.* (2000). Enhanced productivity during controlled proliferation of BHK-21 cells in continuously perfused bioreactors. Biotech. and Bioeng. 69, 266 - 274].

**Fig. 7. Fraction of SEAP expressing clones from stop codon suppression dependent translation constructs after selection.**

BHK-21 cells were transfected using a stop codon suppression dependent *neo* reading frame translating construct (SEAP<sup>stopMneo</sup>) or cotransfection of separate SEAP and *neo* constructs at 1:10

(SEAP + 1/10 *neo*) and 1:1 (SEAP + *neo*) ratios of plasmid DNA, respectively. SEAP activity from drug resistant BHK-21 cell single clones was determined by the agarose overlay technique (Kirchhoff, S. et al. (1995) Identification of mammalian cell clones exhibiting highly regulated expression from inducible promoters. Trends Genet. 11: 219 - 220; McCracken, A. A. et al. (1984) Studies on the secretion of serum proteins from rat hepatoma cells. Hepatology. 4: 715 - 721; Walls, J. D., Grinnell, B. W. (1990) A rapid and versatile method for the detection and isolation of mammalian cell lines secreting recombinant proteins. Biotechniques 8: 138 - 142). The number of SEAP positive staining cells is given relative to the total number of G418 resistant clones obtained.

**Fig. 8: Plasmid map of pSEAPstopneo**

The promoter contains the minimal promoter sequence from the human CytoMegalovirus promoter (CMV), the enhancer elements are derived from the myeloproliferative sarcoma virus promoter, sd-sa are a splice-donor and splice acceptor site from SV40 late splice site, SEAP (545-2062) encodes the secreted form of the placental alkaline phosphatase, dneo (2271-3128) refers to a sequence encoding an N-terminally truncated neo fragment. The locations of these elements are given in the figure in parenthesis.

**Fig. 9: Plasmid map of pSEAPstopMneo**

MSD (2048-2263) encodes a membrane spanning domain derived from the platelet derived growth factor receptor (PDGFR), Mutant refers to the elimination of an non-initiating AUG codon that was present in the original PDGFR membrane spanning domain. The remaining elements are described in the legend of Fig. 8.

**Fig. 10: Plasmid map of pGFPstopneo**

GFP encodes a green fluorescent protein derived from the jellyfish *Aequorea victoria* (Tsien, R. Y. (1998) The green fluorescent protein. Annu. Rev. Biochem. 67: 509 - 544). The remaining elements are described in the legend of Fig. 8.

**Fig. 11: Plasmid map of pGFPstopneoSECIS**

SECIS refers to a selenocystein insertion element that promotes the translation of UGA codons [Walczak, R. et al. (1997) Solution structure of SECIS, the mRNA element required for eukaryotic selenocysteine insertion--interaction studies with the SECIS-binding protein SBP. Biomed. Environ Sci. 10: 177 - 181; Hubert, N. et al. (1996) RNAs mediating cotranslational insertion of selenocysteine in

eukaryotic selenoproteins. Biochimie 78: 590 - 569]. The remaining elements are described in the previous figure legends.

**Fig. 12: DNA-Sequence pGFPstopneo**

All elements are described in the legend of Fig. 10.

**Fig. 13: DNA-Sequence pGFPstopneoSECIS**

All elements are described in the legend of Fig. 11.

**Fig. 14: DNA sequence pSEAPstopMneo**

All elements are described in the legend of Fig. 9.

**Fig. 15: DNA sequence pSEAPstopneo**

All elements are described in the legend of Fig. 8.

**Fig. 16: Influence of selection pressure on the expression levels of producer cells**

Schematic drawing depicting the statistically expected number of producer cells that survive low selection pressure (L) as area below the depicted curve. A fraction of cells with increased expression permits survival even at high selection pressures (H). High selection pressure reduces the screening effort to identify the small number of optimal producer cells (right) by preventing growth of inefficient producer cells (left).

**Fig. 17: Reduced number of surviving clones**

BHK-21 cells were transfected with recombinant plasmid DNA using calcium-phosphate co-precipitation (Wigler M, Silverstein S, Lee LS, Pellicer A, Cheng Y, Axel R. Transfer of purified herpes virus thymidine kinase gene to cultured mouse cells. Cell. 1977 ,11,:223-32 ). Two days after transfection 1mg G418/ml cell culture medium was added and the cells incubated further until no surviving cells could be detected in a control experiment. The bars indicate the fraction of single cell producer clones surviving the selection procedure. The numbers were calculated as average of two independent experiments (SEAP-Stop-M-neo, transfection with DNA of the plasmid pSEAPstopMneo; SEAP + neo, conventional cotransfection with a SEAP coding and neo coding plasmid DNA).

**Fig. 18: Increased frequency of producer clones**

The graph depicts the fraction of single cell colonies that scored positive in a SEAP filter overlay assay (Kirchhoff S, Koster M, Wirth M, Schaper F, Gossen M, Bujard H, Hauser H. Identification of mammalian cell clones exhibiting highly regulated expression from inducible promoters. Trends Genet. 1995;11:219-20. ). The lowest expression level of a cell clone that scored positive in this test was 0,01 pg/ cell. The experimental details are described in the Fig. 17.

**Examples:**

The following examples illustrate the invention, i.e. the feasibility of stop codon suppression-dependent fusion protein translation to improve the coupling of product and selectable marker gene expression.

As shown in Fig. 1, a selectable marker gene was expressed strictly dependent on the translation of a product gene whereby the initiation codon of the selectable marker gene neomycin phosphotransferase (neo) was removed by PCR-mediated mutagenesis thus leading to a truncated neo reading frame which was fused in frame to the complete product gene-reading frame including the termination codon. As examples for product genes the intracellular green fluorescent protein GFP derived from the bioluminescent jellyfish *Aequorea victoria* [Naylor LH. (1999) Reporter gene technology: the future looks bright. *Biochem. Pharmacol.* 58: 749 - 757; Keith, J., Frank, H., Martine, K. (1999) Lowing jellyfish, luminescence and a molecule called coelenterazine *Trends in Biotechnology* 12: 477 - 481; Ellenberg, J., Lippincott-Schwartz, J., Presley, J. F. (1999) Dual-color imaging with GFP variants. *Trends Cell. Biol.* 9: 52 - 56] or a secreted form of human alkaline phosphatase (SEAP) [Berger, J. et al. (1988) The secreted alkaline phosphatase: a powerful new quantitative indicator of gene expression in eukaryotic cells. *Gene* 66: 1 - 10] was used in various configurations (Fig. 1). DNA manipulations were performed according to Maniatis, T. et al. [Maniatis, T., Fritsch, E. F., Sambrook, J., (1982) In: *Molecular cloning. A laboratory manual.* Cold Spring Harbor Laboratory, Cold Spring Harbor, NY]. Manipulations of BHK-21 host cells were performed according to Mueller PP *et al.* or Geserick C *et al.* [Mueller PP *et al.* (1999) Recombinant glycoprotein product quality in proliferation controlled BHK-21 cells. *Biotech. and Bioeng.* 65: 529 - 536 and Geserick, C. et al. (2000) Enhanced productivity during controlled proliferation of BHK-21 cells in continuously perfused bioreactors. *Biotech and Bioeng* 69: 266 - 274].

**Example 1: Improved coupling of gene expression by stop codon suppression-dependent fusion protein translation.**

The GFP<sub>neo</sub> construct encoded a fusion protein with an N-terminal GFP domain and a C-terminal neo domain (Fig. 2A). When the GFP reading frame was terminated by a stop codon (GFP<sub>stopneo</sub>), the encoded GFP protein was expressed from this construct. Ribosomes have a small error rate resulting in C-terminally extended proteins (Fig. 2B). These naturally occurring translational error events were utilized to express minor amounts of GFP<sub>neo</sub> fusion protein (Fig. 2B).

This natural rate of mistranslation generated sufficient neomycin phosphotransferase activity to render transfected BHK-21 cells G418 resistant (see Fig. 3; GFP<sub>stopneo</sub>). The expected GFP-neo fusion protein conferred neo activity and G418 resistance to the transfected cells. A construct (GFP<sub>neo</sub>) without in-frame stop codon at the end of the GFP reading frame was used in a parallel experiment to demonstrate that a GFP-neo fusion protein confers G418 resistance (Geserick, C. et al. (2000) Enhanced

productivity during controlled proliferation of BHK-21 cells in continuously perfused bioreactors. Biotech. and Bioeng. 69: 266 - 274) (see Fig. 3).

In addition, a SECIS sequence has been added to the stop-codon suppression-dependent construct (GFPstopneoSECIS). All described constructs resulted in G418-resistant colonies, whereas mock transfected cells, i.e. cells that were treated identically except that the DNA was omitted in the procedure, remained G418 sensitive. This demonstrated that all constructs lead to the synthesis of sufficient gene products, most likely GFP-neo fusion proteins, with neo activity high enough to confer G418-resistance to transfected cells.

Limiting translation of the neo reading frame led to survival in the presence of G418 of only those transfected cell clones that showed high expression levels. The level of product formation in the resistant cell clones was determined by measuring the GFP fluorescence intensity. For this purpose, a cell suspension in PBS with 5 % FCS was stained with 0.5 microgramm/ml propidium iodide and analyzed with a FACScan, Beckton Dickinson (see Fig. 4). Fluorescence levels in cells transfected with the GFP stop-codon containing construct (see Fig. 4; GFPstopneo) were above those obtained from control constructs (GFPneo, or GFPstopneoSECIS). This demonstrates a limiting activity of the neo gene product that can advantageously be used to isolate high producer cell clones.

Stop codon dependent translational fusion provided a strict dependence of selectable marker gene expression on the expression of the gene of interest. Product formation was maintained even during prolonged growth in the presence of 1mg/ml G418 in the culture medium (see Fig. 5). In contrast, expression of conventionally co-transfected constructs in the same cells has been shown to decrease with time [Geserick, C. et al. (2000) Enhanced productivity during controlled proliferation of BHK-21 cells in continuously perfused bioreactors. Biotech. and Bioeng. 69: 266 - 274].

#### **Example 2: Secreted product protein - stop codon suppression-dependent fusion protein translation**

The selection system was applicable to secreted product proteins, e.g. secreted alkaline phosphatase (SEAP). SEAP activity from stop-codon dependent translated constructs (SEAPstopMneo, for pSEAPstopMneo see Fig. 8) was compared with a control (SEAP) in transient transfection assays. SEAP activity was determined according to Berger, J. et al. [Berger, J. et al. (1988) The secreted alkaline phosphatase: a powerful new quantitative indicator of gene expression in eukaryotic cells. Gene 66: 1 - 10] (see Fig. 6). The data showed that extension of the product mRNA with neo coding sequences did not impede SEAP expression efficiency in the absence of selection pressure. All constructs led to similar SEAP expression, indicating that the added neo sequences had no significant detrimental effect on expression levels.

The fraction of BHK-21 producer cells among clones surviving the selection in the presence of G418 [Geserick, C. et al. (2000) Enhanced productivity during controlled proliferation of BHK-21 cells in continuously perfused bioreactors. *Biotech. and Bioeng.* 69: 266 - 274] was a measure of the stringency of the coupling between product gene expression and selectable marker gene expression. The fraction of producer clones was determined after selection for survival and growth in the presence of G418 (see Fig. 7 and Tab. 1). The number of producer cell clones (Fig. 7 and Tab. 1, SEAPstopMneo) was significantly higher than in the control transfections (Fig. 7 and Tab. 1, SEAP + neo), suggesting that stop codon-dependent translation selection results in a superior coupling rate.

**Table 1.** Efficiency of clone formation from stop codon suppression dependent translation constructs after selection.

BHK-21 cells were transfected by the calcium phosphate coprecipitation method using a stop codon suppression dependent *neo* reading frame translating construct (SEAPstopMneo) or cotransfection of separate SEAP and *neo* constructs at 1:10 (SEAP + 1/10 *neo*) and 1:1 (SEAP + *neo*) ratios of plasmid DNA, respectively. 1000 microgram per ml culture was added and the incubation continued until all cells from a non-transfected control culture were dead. Producer clones were determined by an agarose overlay diffusion filter binding assay for alkaline phosphatase activity.

Construct:	SEAPstopMneo	SEAP+1/10 neo	SEAP+neo
Number of clones	255	27	394
Number of producer clones	93	3	14

### Example 3: Construction of pGFPstopneo

The plasmid pGFPstopneo was constructed from an existing plasmid PMC2LUZI [Mielke, C., Tummler, M., Bode, J. A., simple assay for puromycin N-acetyltransferase: selectable marker and reporter. *Trends Genet.* (1995) 11: 258 -259] by standard recombinant DNA techniques [Sambrook et al. (1989) In: *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, second edition].

The *neo* gene was derived from the bacterial transposon Tn 5. To create neo fusion proteins, the AUG initiation codon was deleted from the *neo* reading frame by PCR-mediated mutagenesis using the primers PMU212 5'-GCGGTCGACGGCCCGGGTCAGAAGAACTC-3' and PMU213 5'-GGCACTAGTCACACCGGTGGATTGCACGCAGGTTCTCCGGCC-3' and a fusion PCR was performed with eGFP (Clontech Laboratories, Inc., Palo Alto, CA.) to generate the GFPstopneo reading frame fusion. The resulting PCR fragment was cleaved using the restriction endonucleases *AscI* and *SmaI* and ligated to the vector backbone in place of the PAC gene downstream of the IRES (Internal Ribosome Entry Site) element after cleavage by the same enzymes. The ligation mixture was

transformed into competent *E. coli* cells (*E. coli* JM109) by electroporation. Transformants were selected on Ampicillin containing solid medium. DNA was isolated from individual colonies and tested for the presence of expected restriction fragments. The critical sequences surrounding the stop codon were confirmed by enzymatic sequencing [Sanger, F., Coulson, A. R. (1975) A rapid method for determining sequences in DNA by primed synthesis with DNA polymerase. *J. Mol. Biol.* 94: 441 - 448].

#### **Example 4: Construction of pGFPstopneoSECIS**

A SECIS consensus element was synthesized as two complementary oligonucleotides of the following sequence (5' to 3') :

SECIS1: 5'-CAGGGCCTTGGATGAGATGAGTATCATAAACTATGACCTAGGGGATACTCATGGATC-AAGGCCTTGGGC-3'

and SECIS2: 5'-CCAAGGCCTTGATCCATGAGTATCCCCTAGGTCATAGTTTATGATACTCATCTCA-TCCAAGGCCCTGGGC-3'.

It was inserted downstream of the *neo* reading frame into the unique SacII restriction site of the plasmid pGFPstopneo by using T4 DNA ligase.

#### **Example 5: Construction of pSEAPstopneo**

pSEAPstopneo is a plasmid vector comprising the secreted alkaline phosphatase-reading frame followed the *neo* reading frame. The construct was derived from pSEAPstopMneo by cleavage with *Bst*BI and religation according to Maniatis, T. et al. [Maniatis, T., Fritsch, E. F., Sambrook, J., (1982) In: *Molecular cloning. A laboratory manual.* Cold Spring Harbor Laboratory, Cold Spring Harbor, NY].

#### **Example 6: Construction of pSEAPstopMneo**

pSEAPstopMneo is a plasmid vector comprising the secreted alkaline phosphatase-reading frame followed by a Stop codon and a membrane-spanning domain and by the *neo* reading frame. The construct was derived from the plasmid pMPSVHE [Artelt, P. et al. (1988) Vectors for efficient expression in mammalian fibroblastoid, myeloid and lymphoid cells via transfection or infection. *Gene* 68: 213 - 219) and the gene encoding the secreted form of alkaline placental phosphatase [Berger, J. et al. (1988) The secreted alkaline phosphatase: a powerful new quantitative indicator of gene expression in eukaryotic cells. *Gene* 66: 1 - 10). The membrane-spanning domain is derived by PCR amplification from the pHook-1(TM) vector (Invitrogen, Carlsbad, CA) using the two primers PMU202 and PMU203 and was inserted as a *Mfe*I restriction fragment into the unique *Eco*RI site of BMS2, such that both sites were destroyed. The initiation codon deficient *neo* reading frame was inserted as a DNA fragment isolated after enzymatic cleavage of pGFPstopneo-SECIS DNA with *Sal*I and *Spe*I, resulting in pSEAPstop(pA)Mneo. This plasmid DNA was then cut with *Bsp*EI such that the p(A) sequence was removed. The remaining plasmid DNA was religated, resulting in pSEAPstopMneo.



PMU202: 5'-CGCCAATTGGTCGACACTAGTTTCGAAACGTGGCTTCTTCTGCCAAAGGATGATGA-GGA-3'

and PMU203:

5'-GCGCAATTGACGCGTTGAACGCGTTCCGGAATAAATAAATCCGGATTGGAACAAAAACTCATCT-CAGAAG-3'

#### **Example 7: Efficient Selection of cell clones highly expressing a gene of interest**

The efficiency of recombinant gene expression after integration of a recombinant gene into the genome of a mammalian cell is dependent on a number of variables. Even if a selection marker is expressed in a way that the recombinant cell survives selection conditions, the expression of the selectable marker gene does not directly correlate with the expression level of the gene of interest. Selection procedures known from state of the art result in a large fraction of low producer cell clones that may express below 1 microgramm recombinant protein per 10 000 000 cells per day and in a small fraction of cell clones highly expressing a gene of interest, i.e. cells that express above 1 microgramm per 10 000 000 cells per day. These cell clones highly expressing a gene of interest can be identified only with a considerable screening effort. The screening effort can be reduced as follows: (1) Increasing the selection pressure to eliminate low producer cells and thus to reduce the number of cell clones that must be screened, and (2) establishing a tight link between the expression of the selectable marker gene and the expression of the gene of interest to avoid the isolation of clones that efficiently express only the selectable marker gene and not the gene of interest. As a consequence, the average expression level of the cell clones surviving selection conditions will be higher (Fig. 16).

To demonstrate that a recombinant gene expression vector as disclosed herein can be used in a process for efficiently selecting a host cell clone highly expressing said gene of interest, pSEAPstopMneo plasmid DNA was transfected into BHK-21 cells using calcium-phosphate co-precipitation (Wigler M, Silverstein S, Lee LS, Pellicer A, Cheng Y, Axel R. Transfer of purified herpes virus thymidine kinase gene to cultured mouse cells. Cell. 1977,11: 223-32). Two days after transfection, selection conditions were applied by adding 1mg/ml G418 to cell culture medium. Cells were incubated under said selection conditions until no living cells could be detected in a control experiment (cells without a selectable marker gene were also incubated in cell culture medium containing 1mg/ml G418). In a parallel experiment, cells were co-transfected with (1) a SEAP encoding plasmid DNA construct and (2) a neo plasmid DNA construct according to a conventional co-transfection procedure. As a result, the number of surviving cell clones obtained with the pSEAPstopMneo plasmid DNA was reduced by at least 50% indicating that pSEAPstopMneo can be used to reduce the screening effort (Fig. 17).

To demonstrate that the average expression rate of the gene of interest is higher in cell clones comprising pSEAPstopMneo, approximately 100 cell clones obtained with the above mentioned experiment were scored in a SEAP filter overlay assay (Kirchhoff S, Koster M, Wirth M, Schaper F, Gossen M, Bujard H, Hauser H. Identification of mammalian cell clones exhibiting highly regulated

expression from inducible promoters. Trends Genet. 1995; 11: 219-20). As a result, 14% of cell clones transfected with pSEAPstopMneo highly expressed the gene of interest which was three times more than the number of highly expressing cell clones obtained from co-transfecting BHK-21 cells with a SEAP encoding plasmid DNA construct and a neo plasmid DNA construct (Fig. 18). This result clearly demonstrates the use of a recombinant gene expression vector - as disclosed herein - in a process to efficiently selecting host cell clones highly expressing a gene of interest.

**Claims:**

1. A recombinant gene expression vector comprising in the following order a promoter sequence, a gene of interest, a translational stop signal and translationally linked to said gene of interest a selectable marker gene whereby a recombinant gene expression vector is excluded which recombinant gene expression vector contains in the following order a promoter sequence, a gene encoding  $\beta$ -galactosidase, a TGA stop codon, and translationally linked to said  $\beta$ -galactosidase gene a gene encoding luciferase.
2. A recombinant gene expression vector as claimed in claim 1, comprising at least two genes of interest.
3. A recombinant gene expression vector according to claim 1, wherein said selectable marker gene encodes a functional protein for selection of host cells comprising said recombinant gene expression vector and expressing said selectable marker gene.
4. A recombinant gene expression vector according to claim 1, wherein said translational stop signal is at least one stop codon selected from the group of TAA, TGA and TAG.
5. A recombinant gene expression vector according to claim 1, comprising downstream of said selectable marker gene a SECIS element which modulates the expression of said selectable marker gene from the corresponding mRNA.
6. A recombinant gene expression vector, comprising in the following order a promoter sequence, a gene of interest, a translational stop signal and translationally linked to said gene of interest a selectable marker gene wherein the gene of interest encoding a secreted product protein and the selectable marker gene are separated by a translational stop signal and an in frame stop transfer sequence.
7. A host cell comprising a recombinant gene expression vector as claimed in claims 1 to 6, which host cell is capable of expressing both said gene of interest and as a fusion protein said gene of interest and said selectable marker gene.
8. A host cell as claimed in claim 7 which is a host cell line and which is stably maintaining the expressing of said gene of interest during cultivation.
9. A process for producing a host cell clone comprising a recombinant gene expression vector as claimed in claims 1 to 6, which process comprises transforming a host cell with said recombinant gene expression vector as claimed in claims 1 to 6 and selecting a host cell clone comprising said recombinant gene expression vector on the expression of said selectable marker gene.

10. A process for producing a protein encoded by a gene of interest, which process comprises the production of a host cell line comprising a recombinant gene expression vector, the cultivation of said host cell line, stably maintaining the expression of said gene of interest, and recovery of the product of said gene of interest, whereby said recombinant gene expression vector comprises in the following order a promoter sequence, a gene of interest, a translational stop signal and translationally linked to said gene of interest a selectable marker gene, and whereby the production of said host cell line comprises the transformation with said recombinant gene expression vector and the selection of a host cell comprising said recombinant gene expression vector on the expression of said selectable marker gene.
11. A process for producing a secreted product protein encoded by a gene of interest, which process comprises the production of a host cell line comprising a recombinant gene expression vector, the cultivation of said host cell line, stably maintaining the expression of said gene of interest, and recovery of said secreted product protein from the cell's surrounding culture medium, whereby said recombinant gene expression vector comprises in the following order a promoter sequence, a gene of interest encoding a secreted product protein, a translational stop signal, an in frame stop transfer sequence and translationally linked to said gene of interest a selectable marker gene, and whereby the production of said host cell line comprises the transformation with said recombinant gene expression vector and the selection of a host cell comprising said recombinant gene expression vector on the expression of said selectable marker gene.
12. Use of a recombinant gene expression vector as claimed in claims 1 to 6 in a process for efficiently selecting a host cell clone highly expressing a gene of interest.
13. A process of producing a host cell line highly expressing a gene of interest, which process comprises a process for producing a host cell clone as claimed in claim 9 and efficiently selecting a host cell clone highly expresses said gene of interest.




Fig. 1



Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6






Fig. 7



**Fig. 8**

**Fig. 9**




Fig. 10




Fig. 11

**Fig. 12: DNA-Sequence pGFPstopneo**

tcgataatga	aagacccac	ctgtaggttt	ggcaagctag	cttaagtaac	gccattttgc	60
aaggcatggg	aaaaatacat	aactgagaat	agagaagttc	agatcaaggt	caggaacaga	120
gaaacaggag	aatatgggcc	aaacaggata	tctgtggtaa	gcagttcctg	ccccgctcag	180
ggccaagaac	agttggaaca	ggagaattgg	gccaaacagg	atatctgtgg	taagcagttc	240
ctgccccgct	cagggccaag	aacagatggg	ccccagatgc	ggccccgccc	tcagcagttt	300
ctagacataa	cttacggtaa	atggccccgc	tggttgaccg	cccaacgacc	cccgccatt	360
gacgtcaata	atgacgtatg	ttcccatagt	aacgccaata	gggactttcc	attgacgtca	420
atgggtggag	tatttacggg	aaactgcccc	cttggcagta	catcaagtgt	atcatatgcc	480
aagtacgccc	cctattgacg	tcaatgacgg	taaatggccc	gcctggcatt	atgccagta	540
catgacctta	tgggactttc	ctacttgcca	gtacatctac	gtattagtca	tcgctattac	600
catggtgatg	cggttttggc	agtacatcaa	tgggcgtgga	tagcggtttg	actcacgggg	660
atttccaagt	ctccacccca	ttgacgtcaa	tgggagtttg	ttttggcacc	aaaatcaacg	720
ggactttcca	aaatgtcgtg	acaactccgc	cccattgacg	caaatgggcg	gtaggcgtgt	780
acggtgggag	gtctatataa	gcagagctcg	tttagtgaac	cgtcagatcg	cctgggagacg	840
ccatccacgc	tgttttgacc	tccatagaag	acaccgggac	cgatccagcc	tcgaggaact	900
ggaaaaccag	aaagtttaact	ggtaagttta	gtctttttgt	cttttatttc	aggtcccggg	960
tcgaatttgcg	gccgcgaatt	cgagctcggg	accatggaag	acgccccaaa	cataaagaaa	1020
ggcccggcgc	cattctatcc	gctagaggat	ggaaccgctg	aagagcaact	gcataaggct	1080
atgaagagat	acgccctggg	tcctgaaaca	attgctttta	cagatgcaca	tatcgagggtg	1140
aacatcacgt	acgcggaata	cttcgaaatg	tccgttcggg	tggcagaagc	tatgaaacga	1200
tatgggctga	atacaaatca	cagaatcgtc	gtatgcagtg	aaaactctct	tcaattcttt	1260
atgccggtgt	tgggcgcggt	atttatcgga	gttgacagttg	cgcccgcgaa	cgacattttat	1320
aatgaacgtg	aattgctcaa	cagtatgaac	atttcgcagc	ctacggtagt	gtttgtttcc	1380
aaaaaggggt	tgcaaaaaat	tttgaacgtg	caaaaaaat	taccaataat	ccagaaaatt	1440
attatcatgg	attctaaaaac	ggattaccag	ggatttcagt	cgatgtacac	gttcgtcaca	1500
tcttatctac	ctcccggttt	taatgaatac	gattttgtac	cagagtcctt	tgactcgtgac	1560
aaaacaattg	cactgataat	gaactcctct	ggatctactg	ggttacctaa	gggtgtggcc	1620
cttcgcata	gaactgcctg	cgtcagattc	tcgcatgcca	gagatcctat	ttttggcaat	1680
caaatcattc	cggatactgc	gatttttaagt	gttggtcaat	tccatcacgg	ttttggaatg	1740
tttactacac	tcggatatatt	gatatgtgaa	tttcgagtcg	tcaaatgtat	agatttgaag	1800
aagagctgtt	ttaacgatcc	cttcaggatt	acaaaattca	aagtgcgttg	cttagtacia	1860
accctatttt	cattcttcgc	caaaagcact	ctgattgaca	aatacgattt	atctaattta	1920
cacgaaattg	cttctggggg	cgcacctctt	tcgaaagaag	tcgggggaagc	ggttgcaaaa	1980
cgcttccatc	ttcaagggat	acgacaagga	tatgggctca	ctgagactac	atcagctatt	2040
ctgattacac	ccgaggggga	tgataaacccg	ggcgcggtcg	gtaaaattgt	tccatttttt	2100
gaagcgaagg	ttgtggatct	ggataccggg	aaaacgctgg	gcgttaatca	gagaggcgaa	2160
ttatgtgtca	gaggacctat	gattatgtcc	ggttatgtaa	acaatccgga	agcgaccaac	2220
gccttgattg	acaaggatgg	atggctacat	tctggagaca	tagcttactg	ggacgaagac	2280
gaacacttct	tcatagttga	ccgcttgaag	tctttaatta	aatacaaagg	ataccagggtg	2340
gccccgcgtg	aattggagtc	gatattgtta	caacacccca	acatcttcga	cgcgggcgtg	2400
gcaggtcttc	ccgacgatga	cgccggtgaa	cttcccgcgc	ccgttggtgt	tttgaggcac	2460
ggaaagacga	tgacggaaaa	agagatcgtg	gattacgtcg	ccagtcaagt	aacaaccgcc	2520
aaaaagttgc	gcggaggagt	tgtgttttg	gacgaagtc	cgaaaggctc	taccggaaaa	2580
ctcgacgcaa	gaaaaatcag	agagatcctc	ataaaggcca	agaaggcgcg	aaagtccaaa	2640
ttgtaaaaatg	taactgtatt	cagcgatgac	gaaattctta	gctattgtaa	tgactctaga	2700
gtcgacctgc	agcccaagct	tttaaaacag	ctctgggggt	gtaccacccc	cagaggccca	2760
cgtggcggct	agtactccgg	tattgcggta	cccttgtagc	cctgttttat	actcccttcc	2820
cgtaacttag	acgcacaaaa	ccaagttcaa	tagaaggggg	tacaaaccag	taccaccacg	2880
aacaagcact	tctgtttccc	cggatgatgtc	gtatagactg	cttgcggtgg	tgaaagcgac	2940
ggatccgtta	tccgcttatg	tacttcgaga	agcccagtac	cacctcgga	tcttcgatgc	3000
gttgcgctca	gcaactcaacc	ccagagtgtg	gcttaggctg	atgagtctgg	acatccctca	3060
ccggtgacgg	tgggtccaggc	tgcgttgggc	gcctacctat	ggctaaccgc	atgggacgct	3120
agttgtgaac	aaagtgtgaa	gagcctattg	agctacataa	gaatcctccg	gcccctgaat	3180
cgggctaate	ccaacctcgg	agcaggtggt	cacaaaccag	tgattggcct	gtcgtaacgc	3240
gcaagtccgt	ggcggaaccg	actacttttg	gtgtccgtgt	ttccttttat	tttattgtgg	3300
ctgcttatgg	tgacaatcac	agattgttat	cataaagcga	attggattgc	ggccgcgcgc	3360
cacgcgtaga	tctacgcgtg	gcgcgccatg	gtgagcaagg	gcgaggagct	gttcaccggg	3420
gtggtgcccc	tcctgggtcg	gctggacggc	gacgtaaacg	gccacaagtt	cagcgtgtcc	3480
ggcgagggcg	agggcgatgc	cacctacggc	aagctgaccc	tgaagtccat	ctgcaccacc	3540
ggcaagctgc	ccgtgccctg	gcccaccctc	gtgaccaccc	tgacctacgg	cgtgcagtg	3600

ttcagccgct	accccgacca	catgaagcag	cacgacttct	tcaagtccgc	catgcccga	3660
ggctacgtcc	aggagcgac	catcttcttc	aaggacgacg	gcaactacaa	gacccgcgcc	3720
gaggtgaagt	tcgagggcga	caccctggtg	aaccgcacgc	agctgaaggg	catcgacttc	3780
aaggaggacg	gcaacatcct	ggctggagta	caactacaac	agccacaacg	tctatatcat	3840
ggccgacaag	cagaagaacg	gcatcaaggt	gaacttcaag	atccgccaca	acatcgagga	3900
cggcagcggtg	cagctcgccg	accactacca	gcagaacacc	cccatcggcg	acggccccgt	3960
gctgctgccc	gacaaccact	acctgagcac	ccagtccgcc	ctgagcaaag	accccaacga	4020
gaagcgcgat	cacatggtcc	tgttgaggtt	cgtgaccgcc	gccgggatca	ctctcggcac	4080
ggacgagctg	tacaagacta	gttgaactag	tattgaacaa	gatggattgc	acgcaggttc	4140
tccggccgct	tgggtggaga	ggctattcgg	ctatgactgg	gcacaacaga	caatcggtcg	4200
ctctgatgcc	gccgtgttcc	ggctgtcagc	gcaggggcgc	ccggttcttt	ttgtcaagac	4260
cgacctgtcc	ggtgccttga	atgaactgca	agacgaggca	gcgcggctat	cgtggctggc	4320
cacgacgggc	gttccttgcg	cagctgtgct	cgacgttgct	actgaagcgg	gaaggagctg	4380
gctgctattg	ggcgaagtgc	cggggcagga	tctcctgtca	tctcaccttg	ctcctgccga	4440
gaaagtatcc	atcatggctg	atgcaatgcg	gcggctgcat	acgcttgatc	cggctacctg	4500
cccattcgac	caccaagcga	aacatcgcat	cgagcgagca	cgtactcgga	tggaaagccg	4560
tcttgctgat	caggatgatc	tggacgaaga	gcatcagggg	ctcgcgccag	ccgaactgtt	4620
cgccaggctc	aaggcgagca	tgcccgacgg	cgaggatctc	gtcgtgacct	atggcgatgc	4680
ctgcttgccg	aatatcatgg	tggaaaatgg	ccgcttttct	ggattcatcg	actgtggccg	4740
gctgggtgtg	gcggaccgct	atcaggacat	agcgttggct	acccgtgata	ttgctgaaga	4800
gcttggcggc	gaatgggctg	accgcttcct	cgtgctttac	ggtatcgccg	ctcccgatcc	4860
gcagcgcac	gccttctatc	gccttcttga	cgagtcttcc	tgaccggggc	cgtacgcacc	4920
ctcgccgcgc	gcttcgccga	ctaccccgcc	acgcgccaca	ccgtcgacct	ggaccgccac	4980
atcgagcggg	tcaccgagct	gcaagaatc	ttcctcacgc	gcgtcgggct	cgacatcggc	5040
aaggtgtggg	tcgcggacga	cggcgccgcg	gtggcggctc	ggaccacgcc	ggagagcgtc	5100
gaagcggggg	cgggtgttcgc	cgagatcggc	ccgcgcacgc	ccgagttgag	cgggtcccgc	5160
ctggccgcgc	agcaacagat	ggaaggcctc	ctggcgccgc	accggcccaa	ggagcccgcg	5220
tggttccttg	ccaccgtcgc	cgtctcgccc	gaccaccagg	gcaagggtct	gggcagcgcc	5280
gtcgtgctcc	ccggagtggg	ggcgcccgag	cgcgcggggg	tgcccgcctt	cctggagacc	5340
tccgcgcccc	gcaacctccc	cttctacgag	cggctcggct	tcaccgtcac	cgccgacgtc	5400
gagtgcggca	aggaccgcgc	gacctggtgc	atgaccgcga	agcccggtgc	ctgagcccg	5460
ccccacgacc	gcagcgcgcc	gaccgaaagc	agcgcacgac	cccatgagct	tcgatccaga	5520
catgataaga	tacattgatg	agtttgagca	aaccacaact	agaatgcagt	gaaaaaaatg	5580
ctttatttgt	gaaatttgtg	atgctattgc	tttatttgtg	accattataa	gctgcaataa	5640
acaagttaac	aacaacaatt	gcattcattt	tatgtttcag	gttcaggggg	aggtgtggga	5700
ggttttttaa	agcaagtaaa	acctctacaa	atgtggtatg	gctgattatg	atcctgcctc	5760
gcgcgtttcg	gtgatgacgg	tgaaaacctc	tgacacatgc	agctcccggg	gacggtcaca	5820
gcttgtctgt	aagcggatgc	cgggagcaga	caagcccgtc	agggcgcgct	agcgggtggt	5880
ggcgggtgtc	ggggcgcgag	catgaccacg	tcacgtagcg	atagcggagt	gtatactggc	5940
ttaactatgc	ggcatcagag	cagattgtac	tgagagtgcg	ccatatgtcg	ggcgcgcttg	6000
ctggcggttt	tccataggct	ccgccccctc	gacgagcatc	acaaaaatcg	acgctcaagt	6060
cagaggtggc	gaagaccgac	aggactataa	agataccagg	cgtttcccc	tggaaagctcc	6120
ctcgtgcgct	ctcctgttcc	gaccctgccc	cttaccggat	acctgtccgc	ctttctccct	6180
tcgggaagcg	tggcgctttc	tcatagctca	cgctgtaggt	atctcagttc	ggtgtaggtc	6240
gttcgctcca	agctgggctg	tgtgcacgaa	cccccgcttc	agcccgaccg	ctgcgcctta	6300
tccggttaact	atcgtcttga	gtccaacctg	gtaagacacg	acttatcgcc	actggcagca	6360
gccactggta	acaggattag	cagagcgagg	tatgtaggcg	gtgctacaga	gttcttgaag	6420
tgggtggccta	actacggcta	cactagaagg	acagtatttg	gtatctgcgc	tctgctgaag	6480
ccagttacct	tcggaaaaag	agttggtagc	tcttgatccg	gaaaacaaac	caccgcgtgt	6540
agcgggtggt	tttttggttg	caagcagcag	attacgcgca	gaaaaaaagg	atctcaagaa	6600
gatacctttga	tcttttctac	ggggcttgac	gctcagtgga	acgaaaactc	acgttaaggg	6660
attttgggtca	tgagattatc	aaaaaggatc	ttcacctaga	tcctttttaa	ttaaaaatga	6720
agttttaaat	caatctaaag	tatatatgag	taaacttggt	ctgacagtta	ccaatgctta	6780
atcagtgagg	cacctatctc	agcgatctgt	ctatttcggt	catccatagt	tgcttgactc	6840
cccgtcgtgt	agataactac	gatacgggag	ggcttaccat	ctggccccag	tgtgcaatg	6900
ataccgcgag	accacgctc	accggctcca	gatttatcag	caataaacca	gccagccgga	6960
agggccgagc	gcagaagtgg	tcctgcaact	ttatccgcct	ccatccagtc	tattaattgt	7020
tgccgggaag	ctagagtaag	tagttcgcca	gttaatatgt	cgcaacgttg	ttgccattgc	7080
tacagccatg	gtggtgtcac	gctcgctgct	tggtatggct	tcattcagct	ccggttccca	7140
acgatcaagg	cgagttacat	gataccccat	gttgtgcaaa	aaagcgggta	gctccttcgg	7200
tcctccgatc	gttgtcagaa	gtaagttggc	cgcagtggtt	tcactcatgg	ttatggcagc	7260
actgcataat	tctcttactg	tcatgccatc	cgtaagatgc	ttttctgtga	ctggtgagta	7320

ctcaaccaag	tcattctgag	aatagtgtat	gcggcgaccg	agttgctctt	gcccggcgctc	7380
aacacgggat	aataccgcgc	cacatagcag	aactttaaaa	gtgctcatca	ttggaaaacg	7440
ttcttcgggg	cgaaaactct	caaggatctt	accgctgttg	agatccagtt	cgatgtaacc	7500
cactcgtgca	cccaactgat	cttcagcatc	ttttactttc	accagcgttt	ctgggtgagc	7560
aaaaacagga	aggcaaaatg	ccgcaaaaaa	gggaataaag	gcgacacgga	aatggtgaat	7620
actcatactc	ttcctttttc	aatattattg	aagcatttat	cagggttatt	gtctcatgag	7680
cggatacata	tttgaatgta	tttagaaaaa	taaacaaata	ggggttccgc	gcacatttcc	7740
ccgaaaagtg	ccacctgacg	tctaagaaac	cattattatc	atgacattaa	cctataaaaa	7800
taggcgtatc	acgaggccct	ttcgtcttca	agaattgggc	gatcgaccaa	ttctcatggt	7860
tgacagctta	tca					7873

**Fig. 13: DNA-Sequence pGFPstopneoSECIS**

tcgataatga	aagacccac	ctgtaggttt	ggcaagctag	cttaagtaac	gccattttgc	60
aaggcatggg	aaaaatacat	aactgagaat	agagaagttc	agatcaaggt	caggaacaga	120
gaaacaggag	aatatgggcc	aaacaggata	tctgtggtaa	gcagttcctg	ccccgctcag	180
ggccaagaac	agttggaaca	ggagaattgg	gccaaacagg	atatctgtgg	taagcagttc	240
ctgccccgct	cagggccaag	aacagatggg	ccccagatgc	ggccccgccc	tcagcagttt	300
ctagacataa	cttacggtaa	atggccccgc	tggttgaccg	cccaacgacc	cccgccatt	360
gacgtcaata	atgacgtatg	ttcccatagt	aacgccaata	gggactttcc	attgacgtca	420
atgggtggag	tatttacggg	aaactgcccc	cttggcagta	catcaagtgt	atcatatgcc	480
aagtacgccc	cctattgacg	tcaatgacgg	taaatggccc	gcctggcatt	atgccagta	540
catgacctta	tgggactttc	ctacttggca	gtacatctac	gtattagtca	tcgctattac	600
catggtgatg	cggtttttgg	agtacatcaa	tgggcgtgga	tagcggtttg	actcacgggg	660
atttccaagt	ctccacccca	ttgacgtcaa	tgggagtttg	ttttggcacc	aaaatcaacg	720
ggactttcca	aaatgtcgtg	acaactccgc	cccattgacg	caaatgggcg	gtaggcgtgt	780
acggtgggag	gtctatataa	gcagagctcg	tttagtgaac	cgtcagatcg	cctgggagacg	840
ccatccacgc	tgttttgacc	tccatagaag	acaccgggac	cgatccagcc	tcgaggaact	900
ggaaaaccag	aaagttaact	ggtaagttaa	gtctttttgt	cttttatttc	aggtcccggg	960
tcgaattgcg	gccgcgaatt	cgagctcggt	accattggaag	acgccaataa	cataaagaaa	1020
ggcccgcgcc	cattctatcc	gctagaggat	ggaaccgctg	aagagcaact	gcataaggct	1080
atgaagagat	acgccctggt	tcctgaaaca	attgctttta	cagatgcaca	tatcgagggtg	1140
aacatcacgt	acgcggaata	cttcgaaatg	tcggttcggt	tggcagaagc	tatgaaacga	1200
tatgggctga	atacaaatca	cagaatcgtc	gtatgcagtg	aaaactctct	tcaattcttt	1260
atgccggtgt	tgggcgcggt	atztatcgga	gttgacagtg	cgcccgcgaa	cgacatttat	1320
aatgaacgtg	aattgctcaa	cagtatgaac	atttcgcagc	ctacggtagt	gtttgtttcc	1380
aaaaaggggt	tgcaaaaaat	tttgaacgtg	caaaaaaaat	taccaataat	ccagaaaatt	1440
attatcatgg	attctaaaac	ggattaccag	ggatttcagt	cgatgtacac	gttcgtcaca	1500
tcttatctac	ctcccggttt	taatgaatac	gattttgtac	cagagtcctt	tgatcgtgac	1560
aaaacaattg	cactgataat	gaactcctct	ggatctactg	ggttacctaa	gggtgtggcc	1620
cttcgcata	gaactgcctg	cgtcagattc	tcgcatgcca	gagatcctat	ttttggcaat	1680
caaatcattc	cggatactgc	gatttttaagt	gttgttcaat	tccatcacgg	ttttggaatg	1740
tttactacac	tcggatatatt	gatattgtgaa	tttcgagtcg	tcaaatgtat	agatttgaag	1800
aagagctgtt	ttaacgatcc	cttcaggatt	acaaaattca	aagtgcgttg	cttagtacia	1860
accctatttt	cattcttcgc	caaaagcact	ctgattgaca	aatacgattt	atctaattta	1920
cacgaaattg	cttctggggg	cgcacctctt	tcgaaagaag	tcgggggaagc	ggttgcaaaa	1980
cgcttccatc	ttcaagggat	acgacaagga	tatgggctca	ctgagactac	atcagctatt	2040
ctgattacac	ccgaggggga	tgataaacgg	ggcgcggtcg	gtaaaattgt	tccatttttt	2100
gaagcgaagg	ttgtggtact	ggataccggg	aaaacgctgg	gcgttaatac	gagaggcgaa	2160
ttatgtgtca	gaggacctat	gattatgtcc	ggttatgtaa	acaatccgga	agcgaccaac	2220
gccttgattg	acaaggatgg	atggctacat	tctggagaca	tagcttactg	ggacgaagac	2280
gaacacttct	tcatagttga	ccgcttgaag	tctttaatta	aatacaaagg	ataccagggtg	2340
gcccccgctg	aattggagtc	gatattgtta	caacacccca	acatcttcga	cgcgggcggtg	2400
gcaggtcttc	ccgacgatga	cgccgggtgaa	cttcccgcgc	ccgttggtgt	tttgaggcac	2460
ggaaagacga	tgacggaaaa	agagatcggt	gattacgtcg	ccagtcaagt	aacaaccgcc	2520
aaaaagttgc	gcggaggagt	tgtgtttgtg	gacgaagtac	cgaaagggtc	taccggaaaa	2580
ctcgacgcaa	gaaaaatcag	agagatcctc	ataaaggcca	agaaggcgcg	aaagtccaaa	2640
ttgtaaaatg	taactgtatt	cagcgatgac	gaaattctta	gctattgtaa	tgactctaga	2700
gtcgacctgc	agcccaagct	tttaaaacag	ctctgggggt	gtaccacccc	cagaggccca	2760
cgtggcggct	agtactccgg	tattgcggta	cccttgtagc	cctgttttat	actcccttcc	2820
cgtaacttag	acgcacaaaa	ccaagttcaa	tagaaggggg	tacaaaccag	taccaccacg	2880
aacaagcact	tctgtttccc	cgggtgatgtc	gtatagactg	cttgcggtgt	tgaaagcgac	2940
ggatccgtta	tccgcttatg	tacttcgaga	agcccagtac	cacctcgga	tcttcgatgc	3000
gttgcgctca	gcactcaacc	ccagagtgtg	gcttaggctg	atgagtctgg	acatccctca	3060
ccggtgacgg	tggtccaggc	tgcgttggcg	gcctacctat	ggctaaccgc	atgggacgct	3120
agttgtgaac	aaagtgtgaa	gagcctattg	agctacataa	gaatccctcc	gcccctgaat	3180
gcggctaate	ccaacctcgg	agcagggtgt	cacaaaccag	tgattggcct	gtcgtaacgc	3240
gcaagtccgt	ggcggaaccg	actactttgg	gtgtccgtgt	ttccttttat	tttattgtgg	3300
ctgcttatgg	tgacaatcac	agattgttat	cataaagcga	attggattgc	ggccgcgcgc	3360
cacgcgtaga	tctacgcgtg	gcgcgccatg	gtgagcaagg	gcgaggagct	gttcaccggg	3420
gtggtgccca	tcctgggtcga	gctggacggc	gacgtaaacg	gccacaagtt	cagcgtgtcc	3480
ggcgagggcg	agggcgatgc	cacctacggc	aagctgaccc	tgaagtcat	ctgcaccacc	3540
ggcaagctgc	ccgtgccctg	gcccaccctc	gtgaccaccc	tgacctacgg	cgtgcagtgc	3600



ttcagccgct	accccgacca	catgaagcag	cacgacttct	tcaagtccgc	catgcccga	3660
ggctacgtcc	aggagcgac	catcttcttc	aaggacgacg	gcaactacaa	gacccgcgcc	3720
gaggtgaagt	tcgagggcga	caccctggtg	aaccgcacgc	agctgaaggg	catcgacttc	3780
aaggaggacg	gcaacatcct	ggctggagta	caactacaac	agccacaacg	tctatatcat	3840
ggccgacaag	cagaagaacg	gcatcaagg	gaacttcaag	atccgccaca	acatcgagga	3900
cggcagcggtg	cagctcgccg	accactacca	gcagaacacc	cccatcgccg	acggccccgt	3960
gctgctgccc	gacaaccact	acctgagcac	ccagtccgcc	ctgagcaaag	accccaacga	4020
gaagcgcgat	cacatggtcc	tgtctggagt	cgtagccgcc	gccgggatca	ctctcggcac	4080
ggacgagctg	tacaagacta	gttgaactag	tattgaacaa	gatggattgc	acgcagggtc	4140
tccggccgct	tgggtggaga	ggctattcgg	ctatgactgg	gcacaacaga	caatcggtcg	4200
ctctgatgcc	gccgtgttcc	ggctgtcagc	gcagggggcg	ccggttcttt	ttgtcaagac	4260
cgacctgtcc	ggtgccttga	atgaactgca	agacgaggca	gcgcggctat	cgtggctggc	4320
cacgacgggc	gttccttgcg	cagctgtgct	cgacgttgct	actgaagcgg	gaaggagctg	4380
gctgctattg	ggcgaagtgc	cggggcagga	tctcctgtca	tctcaccttg	ctcctgcca	4440
gaaagtatcc	atcatggctg	atgcaatgcg	gcggctgcat	acgcttgatc	cggctacctg	4500
cccatcgcac	caccaagcga	aacatcgcat	cgagcgagca	cgtactcgga	tggaaagccg	4560
tcttgtcgat	caggatgatc	tggacgaaga	gcatcagggg	ctcgcgccag	ccgaactgtt	4620
cgccaggctc	aaggcgagca	tgcccagcgg	cgaggatctc	gtcgtgacct	atggcgatgc	4680
ctgcttgccg	aatatcatgg	tggaaaatgg	ccgcttttct	ggattcatcg	actgtggccg	4740
gctgggtgtg	gcggaccgct	atcaggacat	agcgttggct	acccgtgata	ttgttgaaga	4800
gcttggcggc	gaatgggctg	accgcttctc	cgtgctttac	ggtatcgccg	ctcccgatcc	4860
gcagcgcac	gccttctatc	gccttcttga	cgagtcttcc	tgaccggggc	cgtacgcacc	4920
ctgcgcggc	cgcttcgccg	ctaccccgcg	acgcgccaca	ccgtcgacct	ggaccgccac	4980
atcgagcggg	tcaccgagct	gcaagaatcc	ttcctcacgc	gcgtcgggct	cgacatcggc	5040
aaggtgtggg	tcgcggacga	cggcgccgcc	cagggccttg	gatgagatga	gtatcataaa	5100
ctatgacctc	ggggatactc	atggatcaag	gccttggggc	gtggcggctc	ggaccacgcc	5160
ggagagcgtc	gaagcggggg	cgggtgttcg	cgagatcggc	ccgcgcacat	ccgagttgag	5220
cggttcccgc	ctggccgcgc	agcaacagat	ggaaggcctc	ctggcgccgc	accggcccaa	5280
ggagcccgcg	tggttccttg	ccaccgtcgg	cgtctcgccc	gaccaccagg	gcaagggctc	5340
gggcagcgcc	gtcgtgctcc	cggagtgga	ggcgcccgag	cgcgccgggg	tgcccgcctt	5400
cctggagacc	tccgcgcccc	gcaacctccc	cttctacgag	cggctcggct	tcaccgtcac	5460
cgccgagctc	gagtgcccga	aggaccgcgc	gacctggtgc	atgacccgca	agcccggtgc	5520
ctgacgcccc	ccccacgacc	cgcagcgccc	gaccgaaagg	agcgcacgac	cccatgagct	5580
tcgatccaga	catgataaga	tacattgatg	agtttggaca	aaccacaact	agaatgcagt	5640
gaaaaaaatg	ctttattttgt	gaaattttgt	atgctattgc	tttattttgt	accattataa	5700
gctgcaataa	acaagttaac	aacaacaatt	gcattcattt	tatgtttcag	gttcaggggg	5760
aggtgtggga	ggttttttta	agcaagtaaa	acctctacaa	atgtggtatg	gctgattatg	5820
atcctgcctc	gcgcgtttcg	gtgatgacgg	tgaaaacctc	tgacacatgc	agctcccggg	5880
gacggtcaca	gcttgtctgt	aagcggatgc	cgggagcaga	caagcccgtc	agggcgcgct	5940
agcgggtggt	ggcgggtgtc	ggggcgcgac	catgaccagc	tcacgtagcg	atagcggagt	6000
gtatacttgc	ttaactatgc	ggcatcagag	cagattgtac	tgagagtgc	ccatatgtcg	6060
ggccgcggtg	ctggcggttt	tccataggtc	ccgccccctc	gacgagcatc	acaaaaatcg	6120
acgctcaagt	cagaggtggc	gaaacccgac	aggactataa	agataccagg	cgtttcccc	6180
tggaaagctc	ctcgtgcgct	ctcctgttcc	gacctgtccg	cttaccggat	acctgtccgc	6240
ctttctccct	tcgggaagcg	tggcgctttc	tcatagctca	cgctgtaggt	atctcagttc	6300
ggtgtaggtc	gttcgctcca	agctgggctg	tgtgcacgaa	ccccccgttc	agcccagacc	6360
ctgcgcctta	tccggttaact	atcgtcttga	gtccaacctg	gtaagacacg	acttatcgcc	6420
actggcagca	gccactggta	acaggattag	cagagcgagg	tatgtaggcg	gtgctacaga	6480
gttcttgaag	tgggtggccta	actacggcta	cactagaagg	acagtatttg	gtatctgcgc	6540
tctgtgaag	ccagttacct	tccgaaaaag	agttggtagc	tcttgatccg	gcaaacaaac	6600
caccgctggg	agcgttgggt	tttttggttg	caagcagcag	attacgcgca	gaaaaaaagg	6660
atctcaagaa	gatcctttga	tcttttctac	ggggtctgac	gctcagtgga	acgaaaactc	6720
acgttaaggg	attttgggtc	tgagattatc	aaaaaggatc	ttcacctaga	tccttttaaa	6780
ttaaaaatga	agtttttaaa	caatctaaag	tatatatgag	taaacttggt	ctgacagtta	6840
ccaatgctta	atcagtgagg	cacctatctc	agcgatctgt	ctatttcggt	catccatagt	6900
tgcttgactc	cccgctcggt	agataactac	gatacgggag	ggcttaccat	ctggccccag	6960
tgctgcaatg	ataccgcgag	acccacgctc	accggctcca	gatttatcag	caataaacca	7020
gccagccgga	agggccgagc	gcagaagtgg	tcttgcaact	ttatccgcct	ccatccagtc	7080
tattaattgt	tccgggaag	ctagagtaag	tagttcgcca	gttaaatagt	cgcaacgttg	7140
ttgccattgc	tacaggcatc	gtgggtgtcac	gctcgtcggt	tggatgggt	tcattcagct	7200
cgggttccca	acgatcaagg	cgagttacat	gatcccccat	gttgtgcaaa	aaagcgggta	7260
gtccttccg	tctccgatc	gttgtcagaa	gtaagttggc	cgcagtggtta	tcactcatgg	7320

ttatggcagc	actgcataat	tctcttactg	tcatgccatc	cgtaagatgc	ttttctgtga	7380
ctggtgagta	ctcaaccaag	tcattctgag	aatagtgtat	gcggcgaccg	agttgctctt	7440
gcccggcgtc	aacacgggat	aataccgcgc	cacatagcag	aactttaaaa	gtgctcatca	7500
ttggaaaacg	ttcttcgggg	cgaaaactct	caaggatctt	accgctgttg	agatccagtt	7560
cgatgtaacc	cactcgtgca	cccaactgat	cttcagcatc	ttttactttc	accagcgttt	7620
ctgggtgagc	aaaaacagga	aggcaaaatg	ccgcaaaaaa	gggaataagg	gcgacacgga	7680
aatgttgaat	actcatactc	ttcctttttc	aatattattg	aagcatttat	cagggttatt	7740
gtctcatgag	cggatacata	tttgaatgta	tttagaaaaa	taaacaaata	ggggttccgc	7800
gcacatttcc	ccgaaaagtg	ccacctgacg	tctaagaaac	cattattatc	atgacattaa	7860
cctataaaaa	taggcgtatc	acgaggccct	ttcgtcttca	agaattgggc	gatcgaccaa	7920
ttctcatgtt	tgacagctta	tca				7943

**Fig. 14: DNA sequence pSEAPstopMneo**

ctagcttaag	taacgccatt	ttgcaaggca	tgggaaaaat	acataactga	gaatagagaa	60
gttcagatca	aggtcaggaa	cagagaaaca	ggagaatatg	ggccaaacag	gatatctgtg	120
gtaagcagtt	cctgccccgc	tcagggccaa	gaacagttgg	aacaggagaa	ttgggcca	180
caggatatct	gtggtaagca	gttcctgccc	cgctcagggc	caagaacaga	tggtccccag	240
atgcggtccc	gccctcagca	gtttctagag	aacctcaga	tgtttcagg	gtgccccaa	300
gacctgaat	gacctgtgc	cttatttgaa	ctaaccaatc	agttcgcttc	tcgcttctgt	360
tcgcgcgctt	ctgctccccg	agctcaataa	aagagcccac	aacccctcac	tcggcgcgcc	420
agtcctccga	tagactcggt	cgcccgggta	ccgagctcga	attgatctct	cgaggagctg	480
aaaagttaac	tggttaagttt	agtctttttg	tcttttattt	cagggtcccg	atcgaagctt	540
ctgcatgctg	ctgctgctgc	tgctgctggg	cctgaggcta	cagctctccc	tgggcatcat	600
cccagttgag	gaggagaacc	cggacttctg	gaaccgcgag	gcagccgagg	ccctgggtgc	660
cgccaagaag	ctgcagcctg	cacagacagc	cgccaagaac	ctcatcatct	tcctgggcga	720
tgggatgggg	gtgtctacgg	tgacagctgc	caggatccta	aaagggcaga	agaaggacaa	780
actggggcct	gagatacccc	tggccatgga	ccgcttccca	tatgtggctc	tgtccaagac	840
atacaatgta	gacaaacatg	tgccagacag	tggagccaca	gccacggcct	acctgtgcgg	900
ggtcaagggc	aacttccaga	ccattggctt	gagtgcagcc	gcccgttta	accagtgcaa	960
cacgacagc	ggcaacgagg	tcattctcgt	gatgaatcgg	gccaagaaag	cagggaaagt	1020
agtgggagtc	gtaaccacca	cacgagtgcg	gcacgcctcg	ccagccggca	cctacgcccc	1080
cacggtgaac	cgcaactggg	actcggacgc	cgacgtgcct	gcctcggccc	gccaggaggg	1140
gtgccaggac	atcgctacgc	agctcatctc	caacatggac	attgacgtga	tcctaggtgg	1200
aggccgaaag	tacatgtttc	ccatgggaac	cccagaccct	gagtaccag	atgactacag	1260
ccaaggtggg	accaggctgg	acgggaagaa	tctggtgcag	gaatggctgg	cgaagcgcca	1320
gggtgcccgg	tatgtgtgga	accgcaactg	gctcatgcag	gcttccctgg	acccgtctgt	1380
gacccatctc	atgggtctct	ttgagcctgg	agacatgaaa	tacgagatcc	accgagactc	1440
cacactggac	ccctccctga	tggagatgac	agaggctgcc	ctgcgcctgc	tgagcaggaa	1500
cccccgcgcc	ttcttctctc	tcgtggaggg	tggtcgcata	gaccatgggt	atcatgaaag	1560
cagggtttac	cgggcactga	ctgagacgat	catgttcgac	gacgccattg	agagggcggg	1620
ccagctcacc	agcgaggagg	acacgctgag	cctcgtcact	gccgaccact	cccacgtctt	1680
ctccttcgga	ggctaccccc	tgcgaggggg	ctccatcttc	gggctggccc	ctggcaaggc	1740
ccgggacagg	aaggcctaca	cggctctcct	atacggaaac	ggtccagggt	atgtgtctca	1800
ggacggcgcc	cggccggatg	ttaccgagag	cgagagcggg	agccccgagt	atcggcagca	1860
gtcagcagtg	cccctggacg	aagagaccca	cgcaggcgag	gacgtggcgg	tggtcgcgcg	1920
cggcccgcag	gcgcacctgg	ttcacggcgt	gcaggagcag	accttcata	cgcacgtcat	1980
ggccttcgcc	gcctgcctgg	agccctacac	cgcctgcgac	ctggcgcccc	ccgcggcac	2040
caccgacgcc	cgcacccgg	gttgaattga	cgcgtccgga	ttcgaacaaa	aactcatctc	2100
agaagaggat	tggaatgctg	tgggcccagg	cacgcaggag	gtcatcgtgg	tgccacactc	2160
cttgcccttt	aagggtgggtg	tgatctcagc	catcctggcc	ctgggtgggtg	tcaccatcat	2220
ctcccttata	atcctcatca	tcctttggca	gaagaagcca	cgtttcgaaa	ctagtattga	2280
acaagatgga	ttgcacgcag	gttctccggc	cgcttgggtg	gagaggctat	tcggctatga	2340
ctgggcacaa	cagacaatcg	gctgctctga	tgccgcccgt	ttccggctgt	cagcgcaggg	2400
gcgcccgggt	ctttttgtca	agaccgacct	gtccgggtgcc	ctgaatgaac	tgcaagacga	2460
ggcagcgcg	ctatcgtggc	tggccacgac	ggcggttcct	tgcgagctg	tgctcgacgt	2520
tgtcactgaa	gcgggaagg	actggctgct	attgggcgaa	gtgcccgggc	aggatctcct	2580
gtcatctcac	cttgctcctg	cagagaaagt	atccatcatg	gctgatgcaa	tgccggcgct	2640
gcatacgctt	gatccggcta	cctgcccatt	cgaccaccaa	gcgaacatc	gcatcgagcg	2700
agcacgtact	cggatggaag	ccggtcttgt	cgatcaggat	gatctggacg	aagagcatca	2760
ggggctcgcg	ccagccgaac	tggtcgccag	gctcaaggcg	agcatgccc	acggcgagga	2820
tctcgtcgtg	acccatggcg	atgcctgctt	gccgaatatc	atgggtgaaa	atggccgctt	2880
ttctggattc	atcgactgtg	gccggctggg	tgtggcggac	cgctatcagg	acatagcggt	2940
ggctaccggt	gatattgctg	aagagcttgg	cggcgaatgg	gctgaccgct	tcctcggtgt	3000
ttacggtatc	gccgctcccc	attcgcagcg	catcgccttc	tatcgcttc	ttgacgagtt	3060
cttctgacct	gggcccgtac	caccctcgcc	gccgcgttcg	ccgactaccc	cgccacgcgc	3120
cacaccgtcg	accattcccg	atccagacat	gataagatac	attgatgagt	ttggacaac	3180
cacaactaga	attcagtgaa	aaaaatgctt	tatttgtgaa	atttgtgatg	ctattgtctt	3240
atttgttaacc	attataagct	gcaataaaca	agttaacaac	aacaattgca	ttcattttat	3300
gtttcagggt	cagggggagg	tgtgggaggt	tttttaagc	aagtaaaacc	tctacaaatg	3360
tggtatggct	gattatgatc	ctgcctcgcg	cgtttcgggtg	atgacgggtg	aaacctctga	3420
cacatgcagc	tcccggagac	ggtcacagct	tgtctgtgaa	cggatgccgg	gagcagacaa	3480
gcccgtcagg	gcgcgtcagc	gggtgttggc	gggtgtcggg	gcgcagccat	gacccagtc	3540
cgtagcgata	gcggagtgta	tactggctta	actatgcggc	atcagagcag	attgtactga	3600

gagtgcacca	tatcgggccc	cgttgctggc	gtttttccat	aggctccgcc	cccctgacga	3660
gcatcacaaa	aatcgacgct	caagtccagag	gtggcgaaac	ccgacaggac	tataaagata	3720
ccaggcgttt	ccccctggaa	gctccctcgt	gcgctctcct	gttccgaccc	tgccgcttac	3780
cggatacctg	tccgcctttc	tcccttcggg	aagcgtggcg	ctttctcata	gctcacgctg	3840
taggtatctc	agttcgggtg	aggtcgttcg	ctccaagctg	ggctgtgtgc	acgaaccccc	3900
cgttcagccc	gaccgctgcg	ccttatccgg	taactatcgt	cttgagtcca	acccggtaag	3960
acacgactta	tcgccactgg	cagcagccac	tggtaacagg	attagcagag	cgaggatatgt	4020
aggcggtgct	acagagttct	tgaagtgggtg	gcctaactac	ggctacacta	gaaggacagt	4080
atttggtatc	tgcgctctgc	tgaagccagt	taccttcgga	aaaagagttg	gtagctcttg	4140
atccggcaaa	caaaccaccg	ctggtagcgg	tggttttttt	gtttgcaagc	agcagattac	4200
gcgcagaaaa	aaaggatctc	aagaagatcc	tttgatcttt	tctacggggt	ctgacgctca	4260
gtggaacgaa	aactcacggt	aagggatttt	ggatcatgaga	ttatcaaaaa	ggatcttcac	4320
ctagatcctt	ttaaattaaa	aatgaagttt	taaatacaatc	taaagtatat	atgagtaaac	4380
ttggtctgac	agttaccaat	gcttaatcag	tgaggcacct	atctcagcga	tctgtctatt	4440
tcgttcatcc	atagttgcct	gactccccgt	cgtgtagata	actacgatac	gggagggtct	4500
accatctggc	cccagtgtgt	caatgatacc	gcgagaccca	cgctcaccgg	ctccagattt	4560
atcagcaata	aaccagccag	ccggaagggc	cgagcgcaga	agtggctctg	caactttatc	4620
cgcctccatc	cagtctatta	attgttgccg	ggaagctaga	gtaagtagtt	cgccagttaa	4680
tagtttgggc	aacgttgttg	ccattgctac	aggcatcgtg	gtgtcacgct	cgtcgtttgg	4740
tatggcttca	ttcagctccg	gttcccaacg	atcaaggcga	gttacatgat	cccccatgtt	4800
gtgcaaaaaa	gcggttagct	ccttcgggtcc	tccgatcgtt	gtcagaagta	agttggccgc	4860
agtgttatca	ctcatggtta	tggcagcact	gcataattct	cttactgtca	tgccatccgt	4920
aagatgcttt	tctgtgactg	gtgagtactc	aaccaagtca	ttctgagaat	agtgtatgcg	4980
gcgaccgagt	tgctcttgcc	cggcgtcaac	acgggataat	accgcgccac	atagcagaac	5040
tttaaaagtg	ctcatcattg	gaaaacggtc	ttcggggcga	aaactctcaa	ggatcttacc	5100
gctgttgaga	tccagttcga	tgtaacccac	tcgtgcaccc	aactgatctt	cagcatcttt	5160
tactttcacc	agcgtttctg	ggtgagcaaa	aacaggaagg	caaaatgccg	caaaaaaggg	5220
aataagggcg	acacggaaat	gttgaatact	catactcttc	ctttttcaat	attattgaag	5280
catttatcag	ggttattgtc	tcatgagcgg	atacatattt	gaatgtattt	agaaaaataa	5340
acaaataggg	gttccgcgca	catttccccg	aaaagtgcc	cctgacgtat	cgatggtaac	5400
ctctagag						5408

**Fig. 15: DNA sequence pSEAPstopneo**

ctagcttaag	taacgccatt	ttgcaaggca	tgggaaaaat	acataactga	gaatagagaa	60
gttcagatca	aggtcaggaa	cagagaaaca	ggagaatatg	ggccaaacag	gatatctgtg	120
gtaagcagtt	cctgccccgc	tcagggccaa	gaacagttgg	aacaggagaa	ttgggccaaa	180
caggatatct	gtggtaagca	gttcctgccc	cgctcagggc	caagaacaga	tggtccccag	240
atgcggtccc	gccctcagca	gtttctagag	aacctcaga	tgtttccagg	gtgccccaa	300
gacctgaaat	gacctgtgc	cttatattgaa	ctaaccaatc	agttcgcttc	tcgcttctgt	360
tcgcgcgttt	ctgctccccg	agctcaataa	aagagccac	aaccttcac	tcggcgccg	420
agtcctccga	tagactgcgt	cgcccgggta	ccgagctcga	attgatctct	cgaggactg	480
aaaagttaac	tggtaagttt	agtctttttg	tcttttattt	caggctccgg	atcgaagctt	540
ctgcatgctg	ctgctgctgc	tgctgctggg	cctgaggcta	cagctctccc	tgggcatcat	600
cccagttgag	gaggagaacc	cggacttctg	gaaccgcgag	gcagccgagg	ccctgggtgc	660
cgccaagaag	ctgcagcctg	cacagacagc	cgccaagaac	ctcatcatct	tcctgggcga	720
tgggatgggg	gtgtctacgg	tgacagctgc	caggatccta	aaagggcaga	agaaggacaa	780
actggggcct	gagatacccc	tggccatgga	ccgcttccca	tatgtggctc	tgtccaagac	840
atacaatgta	gacaaacatg	tgccagacag	tggagccaca	gccacggcct	acctgtgcgg	900
ggtcaagggc	aacttccaga	ccattggctt	gagtgcagcc	gcccgtttta	accagtgcga	960
cacgacacgc	ggcaacgagg	tcattctccg	gatgaatcgg	gccaagaaag	cagggaagtc	1020
agtgggagtc	gtaaccacca	cacgagtgcg	gcacgcctcg	ccagccggca	cctacgcccc	1080
cacggtgaac	cgcaactggg	actcggacgc	cgacgtgcct	gcctcggccc	gccaggaggg	1140
gtgccaggac	atcgctacgc	agctcatctc	caacatggac	attgacgtga	tcctaggtgg	1200
aggccgaaag	tacatgtttc	ccatgggaac	cccagaccct	gagtaccag	atgactacag	1260
ccaaggtggg	accaggtcgg	acgggaagaa	tctggtgcag	gaatggctgg	cgaagcgcca	1320
gggtgcccgg	tatgtgtgga	accgcaactg	gctcatgcag	gcttccctgg	accggtctgt	1380
gacccatctc	atgggtctct	ttgagcctgg	agacatgaaa	tacgagatcc	accgagactc	1440
cacactggac	ccctccctga	tggagatgac	agaggctgcc	ctgcgcctgc	tgagcaggaa	1500
ccccgcggc	ttcttctctc	tcgtggaggg	tggtcgcata	gacctgggtc	atcatgaaag	1560
cagggtttac	cgggcactga	ctgagacgat	catgttcgac	gacgccattg	agagggcggg	1620
ccagctcacc	agcgaggagg	acacgctgag	cctcgtcact	gccgaccact	cccacgtctt	1680
ctccttcgga	ggctaccccc	tgcgaggggg	ctccatcttc	gggctggccc	ctggcaaggc	1740
ccgggacagg	aaggcctaca	cggctctcct	atacggaaac	gggccagggt	atgtgtctca	1800
ggacggcgcc	cggcgggatg	ttaccgagag	cgagagcggg	agccccgagt	atcggcagca	1860
gtcagcagtg	cccctggacg	aagagaccca	cgcaggcgag	gacgtggcgg	tgttcgcgcg	1920
cggcccgcag	gcgcacctgg	ttcacggcgt	gcaggagcag	accttcatag	cgcacgtcat	1980
ggccttcgcc	gcctgcctgg	agccctacac	cgcctgcgac	ctggcgcccc	ccgcggcac	2040
caccgacgc	gcgcacccgg	gttgaaatga	cgcgtccgga	ataaagatct	tgtcacaga	2100
tctgtgtgtt	ttttccggat	tcgaaactag	tattgaacaa	gatggattgc	acgcagggtt	2160
tccggccgct	tgggtggaga	ggctattcgg	ctatgactgg	gcacaacaga	caatcggtcg	2220
ctctgatgcc	gccgtgttcc	ggctgtcagc	gcaggggcgc	ccggttcttt	ttgtcaagac	2280
cgacctgtcc	ggtgccttga	atgaactgca	agacagggca	gcgcggctat	cgtggctggc	2340
cacgacgggc	gttccttgcg	cagctgtgct	cgacgttgct	actgaagcgg	gaagggactg	2400
gctgctattg	ggcgaagtgc	cggggcagga	tctcctgtca	tctcaccttg	ctcctgccga	2460
gaaagtatcc	atcatggctg	atgcaatgcg	gcggctgcat	acgcttgatc	cggctacctg	2520
cccatttcgac	caccaagcga	aacatcgcat	cgagcgagca	cgtactcgga	tggaaagccg	2580
tcttgtcgat	caggatgatc	tggacgaaga	gcacagggg	ctcgcgccag	cgaagctgtt	2640
cgccaggctc	aaggcgagca	tgcccgcacg	cgaggatctc	gtcgtgacct	atggcgatgc	2700
ctgcttgccg	aatatcatgg	tggaaaatgg	ccgcttttct	ggattcatcg	actgtggccg	2760
gctgggtgtg	gcggaccgct	atcaggacat	agcgttggct	acccgtgata	ttgtgaaga	2820
gcttggcggc	gaatgggctg	accgcttctc	cgtgctttac	ggtatcgccg	ctcccgattc	2880
gcagcgcata	gccttctatc	gccttcttga	cgagttcttc	tgaccggggc	cgtacgcacc	2940
ctcgcgcggc	cgttcgccga	ctaccccgcc	acgcgccaca	ccgtcgacca	attccgatcc	3000
agacatgata	agatacattg	atgagtttgg	acaaaccaca	actagaatgc	agtgaaaaaa	3060
atgctttatt	tgtgaaattt	gtgatgctat	tgttttattt	gtaaccatta	taagctgcaa	3120
taaacaagtt	aacaacaaca	attgcattca	ttttatgttt	caggttcagg	gggaggtgtg	3180
ggaggttttt	taaagcaagt	aaaacctcta	caaatgtggg	atggctgatt	atgatcctgc	3240
ctcgcgcgtt	tcgggtgatga	cgggtgaaaac	ctctgacaca	tgcagctccc	ggagacggtc	3300
acagcttgtc	tgtaaagcga	tgccgggagc	agacaagccc	gtcagggcgc	gtcagcgggt	3360
gttggcgggt	gtcggggcgc	agccatgacc	cagtcacgta	gcgatagcgg	agtgtatact	3420
ggcttaacta	tgcggcatca	gagcagattg	tactgagagt	gcacatatat	gggcccgcgt	3480
gctggcggtt	ttccataggc	tccgcccccc	tgacgagcat	cacaaaaatc	gacgctcaag	3540
tcagaggtgg	cgaaacccca	caggactata	aagataccag	gcgtttcccc	ctggaagctc	3600

cctcgtgcgc	tctcctgttc	cgaccctgcc	gcttaccgga	tacctgtccg	cctttctccc	3660
ttcgggaagc	gtggcgcttt	ctcatagctc	acgctgtagg	tatctcagtt	cgggtgtaggt	3720
cgttcgtctc	aagctgggct	gtgtgcacga	acccccggt	cagcccgacc	gctgcgcctt	3780
atccggtaac	tatcgtcttg	agtccaaccc	ggtaagacac	gacttatcgc	cactggcagc	3840
agccactggg	aacaggatta	gcagagcgag	gtatgtaggc	ggtgctacag	agttcttgaa	3900
gtgggtggcct	aactacggct	acactagaag	gacagtatct	ggtatctgcg	ctctgctgaa	3960
gccagttacc	ttcggaaaaa	gagttggtag	ctcttgatcc	ggcaaacaaa	ccaccgctgg	4020
tagcgggtgg	ttttttgttt	gcaagcagca	gattacgcgc	agaaaaaaag	gatctcaaga	4080
agatcctttg	atctttttcta	cgggggtctga	cgctcagtg	aacgaaaact	cacgttaagg	4140
gattttgggtc	atgagattat	caaaaaggat	cttcacctag	atccttttaa	attaaaaatg	4200
aagttttaaa	tcaatctaaa	gtatatatga	gtaaacttgg	tctgacagtt	accaatgctt	4260
aatcagttag	gcacctatct	cagcgatctg	tctatttctg	tcattccatag	ttgcctgact	4320
ccccgtcgtg	tagataacta	cgatacggga	gggcttacca	tctggcccca	gtgctgcaat	4380
gataccgcga	gaccacgct	caccggctcc	agatttatca	gcaataaacc	agccagccgg	4440
aagggccgag	cgcagaagtg	gtcctgcgaac	tttatccgcc	tccatccagt	ctattaattg	4500
ttgccgggaa	gctagagtaa	gtagtctcgcc	agttaatagt	ttgggcaacg	ttgttgccat	4560
tgctacaggc	atcgtgggtg	cacgctcgtc	gtttggtag	gcttcattca	gctccgggtc	4620
ccaacgatca	aggcgagtta	catgatcccc	catgttggtg	aaaaaagcgg	ttagctcctt	4680
cggtcctccg	atcgttggtc	gaagtaagtt	ggccgcagtg	ttatcactca	tggttatggc	4740
agcactgcat	aattctctta	ctgtcatgcc	atccgtaaga	tgcttttctg	tgactgggtg	4800
gtactcaacc	aagtcattct	gagaatagtg	tatgcggcga	ccgagttgct	cttgcccggc	4860
gtcaacacgg	gataataccg	cgccacatag	cagaacttta	aaagtgtctc	tcattggaaa	4920
acgttcttcg	gggcgaaaac	tctcaaggat	cttaccgctg	ttgagatcca	gttcgatgta	4980
accactcgt	gcaccaact	gatcttcagc	atcttttact	ttcaccagcg	tttctgggtg	5040
agcaaaaaca	ggaaggcaaa	atgccgcgaaa	aaagggaata	agggcgacac	ggaaatgttg	5100
aatactcata	ctcttccttt	ttcaatatta	ttgaagcatt	tatcagggtt	attgtctcat	5160
gagcggatac	atatttgaat	gtatttagaa	aaataaacia	ataggggttc	cgcgcacatt	5220
tccccgaaaa	gtgccacctg	acgtatcgat	ggtaacctct	agag		5264




Fig. 16




Fig. 17




Fig. 18



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/08520

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/67 C12N15/64 C12N15/62 C12N5/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, MEDLINE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01 44516 A (TULARIK INC ;PTC THERAPEUTICS INC (US)) 21 June 2001 (2001-06-21)	1,3,4, 6-8
Y	the whole document	2,5,9-13
X	WO 97 08330 A (CANCER RES CAMPAIGN TECH ;COLLINS MARY KATHARINE LEVINGE (GB); WEI) 6 March 1997 (1997-03-06)	1-4,7-10
Y	cited in the application the whole document	5,6, 11-13
X	ITO W. AND KUROSAWA Y.: "Development of a prokaryotic expression vector that exploits dicistronic gene organization" GENE, vol. 118, 1992, pages 87-91, XP001093736	1,3,4, 6-13
Y	the whole document	2,5
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\* & \* document member of the same patent family

Date of the actual completion of the international search

8 October 2002

Date of mailing of the international search report

24/10/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Armandola, E

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 02/08520

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	IIDA S. ET AL.: "Expression of a downstream gene from a bicistronic transcription unit in transgenic tobacco plants." GENE, vol. 119, 1992, pages 199-205, XP001093735	1,3,4, 6-13
Y	the whole document	2,5
X	ANDRE A ET AL: "Reinitiation of protein synthesis in Escherichia coli can be induced by mRNA cis-elements unrelated to canonical translation initiation signals" FEBS LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 468, no. 1, 18 February 2000 (2000-02-18), pages 73-78, XP004261001 ISSN: 0014-5793	1,3,4,6, 8-10,12, 13
Y	the whole document	2,5,11
X	F. ZINONI ET AL.: "Features of the formate dehydrogenase mRNA necessary for decoding of the UGA codon as selenocysteine" PROC. NATL. ACAD. SCI. USA, vol. 87, June 1990 (1990-06), pages 4660-4664, XP002187233	1,3,4,7, 8
Y	the whole document	2,5,6, 9-13
X	T.M.M SOGAARD ET AL.: "A sensitive assay of translational fidelity (readthrough and termination) in eukaryotic cells" BIOCHEMISTRY (MOSCOW), vol. 64, no. 12, 1999, pages 1408-1417, XP002187234	1,4,6-8
Y	the whole document	2,5,9-13
X	E. GOLDMAN ET AL.: "Efficiencies of translation in three reading frames of unusual non-ORF sequences isolated from phage display" FASEB J., vol. 14, 2000, pages 603-611, XP002187235	1,3,4,7
Y	the whole document	2,5,6, 9-13
	---	
	-/--	

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/08520

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	S. DANIELSEN ET AL.,: "In vitro selection of enzymatically active lipase variants from phage libraries using a mechanism-based inhibitor" GENE, vol. 272, no. 1-2, 11 July 2001 (2001-07-11), pages 267-274, XP004274862	1-4,6-8
Y	* the whole document, in particular page 268, right column, first full-paragraph and page 270, figure 2 *	5,9-13
Y	--- KIRCHHOFF S ET AL: "Identification of mammalian cell clones exhibiting highly regulated expression from inducible promoters" TRENDS IN GENETICS, ELSEVIER, AMSTERDAM, NL, vol. 11, no. 6, June 1995 (1995-06), pages 219-220, XP004207350 ISSN: 0168-9525 the whole document	12,13
A	--- H. KOLLMUS ET AL.,: "Analysis of eukaryotic mRNA structures directing cotranslational incorporation of selenocysteine" NUCLEIC ACID RESEARCH, vol. 24, no. 7, 1996, pages 1195-1201, XP002187237 cited in the application the whole document	1-13
A	--- WO 88 05466 A (CODON) 28 July 1988 (1988-07-28) cited in the application the whole document -----	1-13

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/08520

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0144516	A	21-06-2001	AU 2272001 A EP 1238110 A2 WO 0144516 A2	25-06-2001 11-09-2002 21-06-2001
WO 9708330	A	06-03-1997	AU 6827896 A EP 0846182 A1 WO 9708330 A1 US 6165715 A	19-03-1997 10-06-1998 06-03-1997 26-12-2000
WO 8805466	A	28-07-1988	AU 1187588 A WO 8805466 A2	10-08-1988 28-07-1988